

DAMTP-99-19
 IMAFF-99-FM/08,
 NIKHEF-99-06
 hep-th/9902044

Singular dimensions of the $N = 2$ superconformal algebras II: the twisted $N = 2$ algebra

Matthias Dörrzapf¹ and Beatriz Gato-Rivera^{2,3}

¹*Department of Applied Mathematics and Theoretical Physics
 University of Cambridge, Silver Street
 Cambridge, CB3 9EW, UK*

²*Instituto de Matemáticas y Física Fundamental, CSIC,
 Serrano 123, Madrid 28006, Spain*

³*NIKHEF, Kruislaan 409, NL-1098 SJ Amsterdam, The Netherlands*

ABSTRACT

We introduce a suitable *adapted ordering* for the twisted $N = 2$ superconformal algebra (i.e. with mixed boundary conditions for the fermionic fields). We show that the ordering kernels for complete Verma modules have two elements and the ordering kernels for G -closed Verma modules just one. Therefore, spaces of singular vectors may be two-dimensional for complete Verma modules whilst for G -closed Verma modules they can only be one-dimensional. We give all singular vectors for the levels $\frac{1}{2}$, 1, and $\frac{3}{2}$ for both complete Verma modules and G -closed Verma modules. We also give explicit examples of degenerate cases with two-dimensional singular vector spaces in complete Verma modules. General expressions are conjectured for the relevant terms of all (primitive) singular vectors, i.e. for the coefficients with respect to the ordering kernel. These expressions allow to identify all degenerate cases as well as all G -closed singular vectors. They also lead to the discovery of subsingular vectors for the twisted $N = 2$ superconformal algebra. Explicit examples of these subsingular vectors are given for the levels $\frac{1}{2}$, 1, and $\frac{3}{2}$. Finally, the multiplication rules for singular vector operators are derived using the ordering kernel coefficients. This sets the basis for the analysis of the twisted $N = 2$ embedding diagrams.

1 Introduction

The spectacular gain of importance of superstring theory in physics over the past fifteen years asked for solutions to challenging problems on the mathematics side. Just like for any quantum field theory, a key problem is the study of the underlying symmetry algebra in order to understand the space of states of the physical theory. Almost at the same time as Ademollo et al.¹ identified for the first time the symmetry algebras of the superstrings, Kac was already studying these objects among his classification of superalgebras²⁴. These symmetry algebras are known as superconformal algebras because the Virasoro algebra is a subalgebra due to the conformal invariance on the string world-sheet. The study of the representations of the superconformal algebras is one of these challenging problems for mathematics. Despite its importance, satisfactory answers are still not known for most superconformal algebras. Increasing the number of fermionic currents N the structure of the representations of the superconformal algebras gets more and more complicated. The case of the $N = 1$ superconformal algebra is well understood thanks to many authors^{2, 6, 5, 18, 19, 23, 25, 27, 28, 30}. However, its representation theory is very close to the representation theory of the Virasoro algebra. Already for the $N = 2$ superconformal algebras there are still many open questions, even though much progress has been made in recent years^{8, 12, 11, 20, 21, 14}. For the $N = 2$ superconformal algebras many aspects arise that are new to representations of chiral algebras. One of these aspects is the existence of degenerate singular vectors^{12, 21}. That is, $N = 2$ superconformal Verma modules can have at the same level and with the same charge more than one linearly independent singular vectors. Thus the study of the embedding structure as well as the computation of character formulae for the representations is much more complicated than in the case where the comparison of level and charge is already enough in order to decide if two singular vectors are proportional, as it is for the Virasoro algebra and for the $N = 1$ superconformal algebra.

In physics, the different types of superconformal symmetry algebras, for the same number of fermionic fields N , arise through the different choices of periodicity conditions for the fermionic currents around a closed superstring world-sheet, as well as for different choices of the Virasoro generators (i.e. of the stress-energy tensor). For the standard choice of Virasoro generators, the superconformal algebras resulting from antiperiodic boundary conditions are called Neveu-Schwarz algebras, those corresponding to periodic boundary conditions are called Ramond algebras and those with mixed boundary conditions are called twisted superconformal algebras. The $N = 2$ superconformal algebras fall under four types: the Neveu-Schwarz $N = 2$ algebra, the Ramond $N = 2$ algebra, the topological $N = 2$ algebra, and the twisted $N = 2$ algebra. The first three algebras are isomorphic, nevertheless their highest weight representations are quite different^a. The topological $N = 2$ algebra, which is the symmetry algebra of topological conformal field theory, also plays an important rôle in string theories^{9, 22, 7}. This algebra is also known as the ‘twisted topological’ $N = 2$ algebra because it can be obtained from the Neveu-Schwarz $N = 2$ algebra by modifying the stress-energy tensor by adding the derivative of the $U(1)$ current, procedure known as ‘topological twist’.

We have studied the singular dimensions of the three isomorphic algebras in a recent publication¹⁶, where we presented a formalism - the adapted ordering method - that allows us to compute upper limits for the dimensions of singular subspaces simply by ordering the algebra generators appropriately. If this order is chosen in a suitable way, then the computed upper limits for these singular dimensions may actually be maxima. We applied this method directly to the topological $N = 2$ algebra proving its usefulness, and then we deduced the maximal singular dimensions corresponding to the Neveu-Schwarz and to the Ramond $N = 2$ algebras.

^aWe thank V. Kac for discussions on this point

So far, not much attention has been given to the twisted superconformal algebras, which appear for $N = 2$ for the first time. This is mainly due to the fact that field theories with mixed periodicity conditions have not been considered for string theories. In the $N = 2$ case there is just one twisted algebra whilst for bigger N there are several possible ways of mixing the periodicity conditions of the fermionic fields and thus different twisted superconformal algebras can be found. In this paper we will focus on the twisted $N = 2$ superconformal algebra. We will show that the method of adapted orderings can also be used for this algebra. For the three isomorphic $N = 2$ algebras as well as for the Virasoro algebra we saw¹⁶ that the suitably chosen adapted ordering assigns a special rôle to the powers of the Virasoro generator L_{-1} . Surprisingly enough, in the twisted $N = 2$ case this rôle is taken over for the first time by another generator. Furthermore, we will see that the twisted $N = 2$ algebra requires its two fermionic fields to be mixed rather than kept separate as for the other $N = 2$ algebras. Hence the twisted $N = 2$ algebra behaves very differently from the other $N = 2$ algebras with respect to the adapted ordering method. Nevertheless, the method still works and we can prove that the computed upper limits for the singular dimensions are actually maxima. Even though the twisted $N = 2$ superconformal algebra contains two fermionic fields, at the algebraic level it looks as if it really contains only one. Furthermore the parameter space of the conformal weights is just one-dimensional, unlike the corresponding parameter spaces for the three isomorphic $N = 2$ algebras. For both reasons, one would have naively suggested that the singular dimensions have to be less than or equal to 1. Surprisingly enough we will show that the maximal singular dimension is in fact 2. Surely, these two-dimensional spaces are not spanned by any tangent space of vanishing surfaces corresponding to singular vectors - as it is the case for the three isomorphic $N = 2$ algebras - simply due to the fact that the parameter space is only one-dimensional. The singular dimension 2 arises rather in a completely new way due to the intersection of two different one-parameter families of singular vectors. This is another reason for which the twisted $N = 2$ algebra is so far unique of its kind.

Superconformal algebras seem to have many surprising features. Not only does this paper show that recent achievements follow the correct way to tackle these and to learn much, which will be helpful to analyse even more complicated conformal structures, it also shows that the twisted $N = 2$ algebra gives more new and different insight in superconformal representation theory. Furthermore this paper creates the basis for the study of the embedding structure of the twisted $N = 2$ highest weight representations since we deduce the singular dimensions as well as the multiplication rules for singular vector operators. Both are crucial for the analysis of the embedding structure. We will also give answers to the question whether the twisted $N = 2$ superconformal algebra contains subsingular vectors. The study of the embedding diagrams, that is the structure of descendant (secondary) singular vectors, will be considered in a forthcoming publication.

This paper is organised in the following way. After a small introduction with the necessary facts about the twisted $N = 2$ algebra in section 2, we review the concept of adapted orderings and its main implications in section 3, in a version suitable for the twisted $N = 2$ algebra. In section 4 we then introduce an adapted ordering for the Verma modules of the twisted $N = 2$ algebra. This allows us to compute the upper limits for the singular dimensions and the multiplication rules for singular vector operators in section 5. That these upper limits are in fact maximal dimensions is demonstrated in section 6 using explicit examples which are supplemented by more examples in appendix A. Based on both theoretical results and explicit computational results we give a reliable conjecture for the coefficients of the relevant terms of all twisted $N = 2$ singular vectors in section 7. Using this conjecture we can characterize all cases of degenerate (i.e. two-dimensional) singular vectors for all levels. This conjecture leads to the discovery of subsingular vectors, which are singular in the G -closed Verma modules considered

in section 8 and in appendix B. It also leads to the identification of all G -closed singular vectors, which are the issue of section 9. We conclude the paper with some final remarks in section 10.

2 The twisted $N = 2$ superconformal algebra

The twisted $N = 2$ superconformal algebra J_2 consists of the Virasoro algebra generators^b L_m , $m \in \mathbb{Z}$, corresponding to the stress-energy tensor, a Heisenberg algebra T_r , with half-integral $r \in \mathbb{Z}_{\frac{1}{2}}$, corresponding to the $U(1)$ current, and the fermionic generators G_k , $k \in \mathbb{Z}^{1/2}$, which are the modes of the two spin-3/2 fermionic fields. J_2 satisfies the following commutation relations^c.

$$\begin{aligned}
[L_m, L_n] &= (m - n)L_{m+n} + \frac{C}{12}(m^3 - m)\delta_{m+n,0} , \\
[L_m, G_k] &= \left(\frac{1}{2}m - k\right)G_{m+k} , \\
[L_m, T_r] &= -rT_{m+r} , \\
[T_r, T_s] &= \frac{1}{3}Cr\delta_{r+s,0} , \\
[T_r, G_k] &= G_{r+k} , \\
\{G_k, G_l\} &= 2\delta_{k+l}^{\mathbb{Z}}(-1)^{2k}L_{k+l} - \delta_{k+l}^{\mathbb{Z}_{\frac{1}{2}}}(-1)^{2k}(k - l)T_{k+l} + \frac{C}{3}\delta_{k+l}^{\mathbb{Z}}(-1)^{2k}(k^2 - \frac{1}{4})\delta_{k+l,0} , \\
[L_m, C] &= [T_r, C] = [G_k, C] = 0 ,
\end{aligned} \tag{1}$$

with $\delta_m^S = 1$ if $m \in S$ and $\delta_m^S = 0$ otherwise, $(-1)^{2k} = +1$ for $k \in \mathbb{Z}$ and $(-1)^{2k} = -1$ for $k \in \mathbb{Z}_{\frac{1}{2}}$, $m, n \in \mathbb{Z}$, $r, s \in \mathbb{Z}_{\frac{1}{2}}$, and finally $k, l \in \mathbb{Z}^{1/2}$. For the Neveu-Schwarz, Ramond and topological $N = 2$ algebras one usually chooses a basis in which the odd generators are nilpotent. If we want to keep the L_0 -grading of the basis for the algebra generators then such a choice does not exist for the twisted $N = 2$ algebra. Nevertheless, the squares of the fermionic operators can be expressed in terms of Virasoro operators, as one can deduce easily from the commutation relations Eqs. (1) :

$$G_k^2 = (-1)^{2k}L_{2k} - \delta_{k,0}\frac{C}{24} , k \in \mathbb{Z}^{1/2} . \tag{2}$$

The central term C commutes with all other operators and can therefore be fixed as $c \in \mathbb{C}$. $\mathcal{H}_{J_2} = \text{span}\{L_0, C\}$ defines a commuting subalgebra of J_2 , which can therefore be diagonalised simultaneously. Generators with positive index span the set of *positive operators* J_2^+ of J_2 and likewise generators with negative index span the set of *negative operators* J_2^- of J_2 :

$$J_2^+ = \text{span}\{L_m, T_r, G_k : m \in \mathbb{N}, r \in \mathbb{N}_{\frac{1}{2}}, k \in \mathbb{N}^{1/2}\} , \tag{3}$$

$$J_2^- = \text{span}\{L_{-m}, T_{-r}, G_{-k} : m \in \mathbb{N}, r \in \mathbb{N}_{\frac{1}{2}}, k \in \mathbb{N}^{1/2}\} . \tag{4}$$

^bWe use the notation $\mathbb{N} = \{1, 2, 3, \dots\}$, $\mathbb{N}_0 = \{0, 1, 2, \dots\}$, $\mathbb{N}_{\frac{1}{2}} = \{\frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \dots\}$, $\mathbb{N}^{1/2} = \mathbb{N} \cup \mathbb{N}_{\frac{1}{2}}$, $\mathbb{N}_0^{1/2} = \mathbb{N}_0 \cup \mathbb{N}_{\frac{1}{2}}$, and also $\mathbb{Z} = \{\dots, -1, 0, 1, 2, \dots\}$, $\mathbb{Z}_{\frac{1}{2}} = \{\dots, -\frac{3}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{3}{2}, \dots\}$, $\mathbb{Z}^{1/2} = \mathbb{Z} \cup \mathbb{Z}_{\frac{1}{2}}$.

^cThe earlier way of writing the twisted $N = 2$ commutation relations, as for instance given by Boucher, Friedan, and Kent⁸, is to distinguish the modes of the two fermionic fields explicitly using superscripts 1 and 2. However, algebraically it is not necessary to distinguish them as one of the fields has modes with integral indices and the other one has modes with half-integral indices. In order to avoid the imaginary unit i appearing in the commutation relations, we have, for convenience, redefined iG_r as G_r for $r \in \mathbb{Z}_{\frac{1}{2}}$ compared to the notation in Ref. 8.

The *zero modes* are spanned by $J_2^0 = \text{span}\{L_0, G_0, C\}$ such that the generator G_0 classifies the different types of Verma modules as we will analyse shortly. One usually extends the algebra by the parity operator^d $(-1)^F$ which commutes with all operators L_m , T_r , and C and anticommutes with G_k :

$$\begin{aligned} [(-1)^F, L_m] &= [(-1)^F, T_r] = [(-1)^F, C] = 0, \quad m \in \mathbb{Z}, r \in \mathbb{Z}_{\frac{1}{2}}, \\ \{(-1)^F, G_k\} &= 0, \quad k \in \mathbb{Z}^{1/2}. \end{aligned}$$

The operator $(-1)^F$ will serve us later to distinguish fermionic from bosonic states in the space of states. As $(-1)^F$ commutes with $\mathcal{H}_{J_2}$ it can also be diagonalised simultaneously to $\mathcal{H}_{J_2}$.

As usual, a simultaneous eigenvector $|\Delta\rangle$ of $\mathcal{H}_{J_2}$ and $(-1)^F$ with L_0 -eigenvalue Δ (the conformal weight), C -eigenvalue^e c (the conformal anomaly), and vanishing J_2^+ action is called a *highest weight vector*, corresponding in fact to the state with *lowest* conformal weight in a given representation of the algebra. Unless otherwise stated, we set $|\Delta\rangle$ to have $(-1)^F$ -eigenvalue^f (parity) $+$. Additional zero-mode vanishing conditions are possible only with respect to the operator G_0 which may or may not annihilate a highest weight vector $|\Delta\rangle$ (c.f. Ref. 16, definition 3.A). If G_0 annihilates the highest weight vector we shall denote it by a superscript in $|\Delta\rangle^G$ and call this highest weight vector *G-closed*. Since $G_0^2 = L_0 - \frac{c}{24}$, any G -closed vector necessarily has conformal weight $\Delta = \frac{c}{24}$. Verma modules are defined in the usual way as the left module obtained by acting with the universal enveloping algebra $U(J_2)$ on a highest weight vector:

$$\mathcal{V}_\Delta = U(J_2) \otimes_{\mathcal{H}_{J_2} \oplus J_2^+} |\Delta\rangle, \quad (5)$$

$$\mathcal{V}_{\frac{c}{24}}^G = U(J_2) \otimes_{\mathcal{H}_{J_2} \oplus J_2^+ \oplus \text{span}\{G_0\}} \left| \frac{c}{24} \right\rangle^G, \quad (6)$$

If the highest weight vector is G -closed, then we call the Verma module $\mathcal{V}_{\frac{c}{24}}^G$ (which is not complete therefore) also G -closed.

For an eigenvector Ψ_l^p of $\mathcal{H}_{J_2}$ in $\mathcal{V}_\Delta$ the conformal weight is $\Delta + l$ and the parity is p with $l \in \mathbb{N}_0^{1/2}$ and $p \in \{\pm\}$. For convenience we will sometimes denote the *level* and the *parity* as $|\Psi_l^p|_L = l$ and $|\Psi_l^p|_F = p$. *Singular vectors* $\Psi_l^\pm$ of a Verma module $\mathcal{V}_\Delta$ are eigenvectors of $\mathcal{H}_{J_2}$ with parity $\pm$ that are not proportional to the highest weight vector but are annihilated by J_2^+ . They correspond therefore to the states with lowest conformal weight in a given subrepresentation of the algebra. They may also satisfy an additional vanishing condition with respect to the operator G_0 in which case we add a superscript G in the notation $\Psi_l^{\pm G}$ and call it a *G-closed singular vector*. Obviously G -closed singular vectors satisfy similar restrictions on their conformal weight as G -closed highest weight vectors: $\Delta + l = \frac{c}{24}$. Most singular vectors can be constructed by acting with algebra generators on another singular vector at a lower (or equal) level. These are called *secondary* singular vectors. Otherwise, they are called *primitive* singular vectors. The *singular vector operator* $\theta_l^\pm$ of $\Psi_l^\pm$ is the unique operator in $U(J_2^- \oplus \{G_0\})$ such that $\Psi_l^\pm = \theta_l^\pm |\Delta\rangle$ and similarly for $\Psi_l^{\pm G}$ where $U(J_2^- \oplus \{G_0\})$ is replaced by $U(J_2^-)$.

Singular vectors and all their descendant vectors (i.e. all the states that are produced by acting with algebra generators on the singular vectors) have vanishing (*pseudo*-)norms under the standard inner product. Vectors with zero norm are usually called *null vectors*^g. The quotient

^dNote that due to Eq. (2) the fermion number F is not well defined, however, the *parity* $(-1)^F$ is.

^eFor simplicity we will suppress the eigenvalue of C in $|\Delta, c\rangle$ and simply write $|\Delta\rangle$.

^fInstead of parity ± 1 we shall simply say parity $\pm$.

^gStrictly speaking, *null-vectors* should only be vectors in the kernel of the inner product matrix and hence decouple completely from the whole space of states. If the pseudo-norm is not positive semi-definite there may be vectors with vanishing pseudo-norm that are not null-vectors¹⁵.

module where all null vectors are set to 0 is hence an irreducible highest weight representation. Conversely, all irreducible highest weight representations can be constructed in this way. However, singular vectors and their descendants do not necessarily span the whole submodule of null vectors. The quotient module of the Verma module divided by the submodule spanned by all singular vectors may again contain new singular vectors, called subsingular vectors. Subsingular vectors and their descendants are also null vectors. Subsingular vectors do not appear for the Virasoro algebra neither for the $N = 1$ superconformal algebras. However, they have been discovered for the three isomorphic $N = 2$ superconformal algebras in Refs. 21, 20. In these cases, the existence of subsingular vectors is very closely related to the existence of singular operators annihilating a singular vector as shall be discussed in a forthcoming publication¹⁵. That the twisted $N = 2$ superconformal algebra also has subsingular vectors will be shown in section 8. of this paper.

Even though the set of null vectors consists of singular vectors, subsingular vectors, and all their descendants, the null vectors with lowest level in a Verma module are certainly singular vectors (i.e. annihilated by all the positive operators). Therefore, the lowest level at which the determinant of the inner product matrix vanishes indicates the presence of (at least) one singular vector. Hence the determinant formula is one of the most important tools for analysing highest weight representations. For the twisted $N = 2$ superconformal algebra the determinant formula has been given by Boucher, Friedan, and Kent⁸. For level zero the determinant formulae are $\det M_0^+ = 1$ and $\det M_0^- = \Delta - \frac{c}{24}$. For level $l \in \mathbb{N}^{1/2}$ and parity $\pm$ the determinant formulae are:

$$\det M_l^\pm = \left(\Delta - \frac{c}{24}\right)^{\frac{P(l)}{2}} \prod_{\substack{1 \leq r, s \leq 2l \\ s \text{ odd}}} \left[2\left(\frac{c}{3} - 1\right)\left(\Delta - \frac{c}{24}\right) + \frac{1}{4}\left(\left(\frac{c}{3} - 1\right)r + s\right)^2\right]^{P(l - \frac{rs}{2})}, \quad (7)$$

where r and s are positive integers (s odd). The partition function $P(l)$, that can be found in Ref. 8, satisfies $P(0) = 1$. The two parity sectors $\pm$ have exactly the same determinant expressions. This correspondence is in fact even closer since the operator G_0 interpolates between the two sectors, provided G_0 does not act on any G -closed vector. Let us analyse this correspondence closer for singular vectors.

For a given singular vector $\Psi_l^+ = \theta_l^+ |\Delta\rangle \in \mathcal{V}_\Delta$ with singular operator θ_l^+ we can construct two negative parity singular vectors:

$$\Psi_{G,l}^- = \theta_l^+ G_0 |\Delta\rangle, \quad (8)$$

$${}_G\Psi_l^- = G_0 \theta_l^+ |\Delta\rangle. \quad (9)$$

Likewise, G_0 constructs two positive parity singular vectors from a singular vector $\Psi_l^- = \theta_l^- |\Delta\rangle \in \mathcal{V}_\Delta$:

$$\Psi_{G,l}^+ = \theta_l^- G_0 |\Delta\rangle, \quad (10)$$

$${}_G\Psi_l^+ = G_0 \theta_l^- |\Delta\rangle. \quad (11)$$

One might therefore believe that singular vectors always come in groups of (at least) 4. However, this assumes that these 4 vectors are actually linearly independent, what is not true as one can guess from the fact that $P(0) = 1$ in the determinant formulae, pointing towards the existence of, generically, one singular vector of each parity at the same level in the same Verma module. This issue will be considered in section 7. In the case of G -closed Verma modules or G -closed singular vectors some of these vectors are actually trivial, an issue that is important for the discussion of embedding diagrams.

For elements Y of J_2 which are eigenvectors of $\mathcal{H}_{J_2}$ and $(-1)^F$ with respect to the adjoint representation we define the level $|Y|_L$ as $[L_0, Y] = |Y|_L Y$ and in addition the parity $|Y|_F$ as $[(-1)^F, Y] = |Y|_F Y$. In particular, elements of the form

$$Y = L_{-m_L} \dots L_{-m_1} T_{-s_T} \dots T_{-s_1} G_{-k_G} \dots G_{-k_1} T_{-\frac{n}{2}}^{r_1} G_{-\frac{1}{2}}^{r_1} G_0^{r_2}, \quad (12)$$

where $r_1, r_2 \in \{0, 1\}$ and any reorderings of Y have level $|Y|_L = \sum_{j=1}^L m_j + \sum_{j=1}^T s_j + \sum_{j=1}^G k_j + \frac{n+r_1}{2}$ and parity $|Y|_F = (-1)^{G+r_1+r_2}$. For these elements we shall also define their *length* $\|Y\| = L + T + G$. For the trivial case we set $|1|_L = \|1\| = 0$ and $|1|_F = +1$. For convenience we define the following sets of negative operators for $l \in \mathbb{N}^{1/2}$

$$\mathbf{L}_l = \{Y = L_{-m_L} \dots L_{-m_1} : m_L \geq \dots \geq m_1 \geq 1, |Y|_L = l\}, \quad (13)$$

$$\mathbf{T}_l = \{Y = T_{-s_T} \dots T_{-s_1} : s_T \geq \dots \geq s_1 \geq \frac{3}{2}, |Y|_L = l\}, \quad (14)$$

$$\mathbf{G}_l = \{Y = G_{-k_G} \dots G_{-k_1} : k_G > \dots > k_1 \geq 1, |Y|_L = l\}, \quad (15)$$

$$\mathbf{L}_0 = \mathbf{T}_0 = \mathbf{G}_0 = \{1\}. \quad (16)$$

We can now define a graded basis for the Verma modules which we use as the standard basis and on which we shall later construct the adapted ordering. A preliminary step is to define for $l \in \mathbb{N}_0^{1/2}$

$$\begin{aligned} \mathcal{S}_l^\pm &= \{Y = LTG : L \in \mathbf{L}_m, T \in \mathbf{T}_r, G \in \mathbf{G}_k, \\ &\quad |Y|_L = l = m + r + k, |Y|_F = \pm 1 = |G|_F, m, r, k \in \mathbb{N}_0^{1/2}\}. \end{aligned} \quad (17)$$

This leads for $l \in \mathbb{N}_0^{1/2}$ to:

$$\begin{aligned} \mathcal{C}_l^\pm &= \left\{ S_{k,p} T_{-\frac{1}{2}}^{2l-2k-r_1} G_{-\frac{1}{2}}^{r_1} G_0^{r_2} : S_{k,p} \in \mathcal{S}_k^p, k \in \mathbb{N}_0^{1/2}, r_1, r_2 \in \{0, 1\}, \right. \\ &\quad \left. 2l - 2k - r_1 \geq 0, p(-1)^{r_1+r_2} = \pm 1 \right\}. \end{aligned} \quad (18)$$

A typical element of $\mathcal{C}_l^\pm$ is hence of the form

$$Y = L_{-m_L} \dots L_{-m_1} T_{-s_T} \dots T_{-s_1} G_{-k_G} \dots G_{-k_1} T_{-\frac{n}{2}}^{r_1} G_{-\frac{1}{2}}^{r_1} G_0^{r_2}, \quad (19)$$

$r_1, r_2 \in \{0, 1\}$, such that $|Y|_L = m$ and $|Y|_F = \pm 1$. $S_{k,p}$ of $Y \in \mathcal{C}_l^\pm$ is called the *leading part* of Y and is denoted by Y^* . Hence, we can define the following standard basis:

$$\mathcal{B}_\Delta = \left\{ X |\Delta\rangle : X \in \mathcal{C}_l^\pm, l \in \mathbb{N}_0^{1/2} \right\}, \quad (20)$$

obtaining finally the Verma module $\mathcal{V}_\Delta = \text{span}\{\mathcal{B}_\Delta\}$.

The basis Eq. (20) is naturally $\mathbb{N}_0^{1/2} \otimes \{\pm 1\}$ graded with respect to their L_0 and $(-1)^F$ eigenvalues, where the L_0 -eigenvalue (the level) is seen relative to the eigenvalue Δ of the highest weight vector.

The *normal form* of an eigenvector of the commuting subalgebra $\mathcal{H}_{J_2}$ is defined as the basis decomposition with respect to the standard basis Eq. (20).

$$\Psi_l^p = \sum_{X \in \mathcal{C}_l^p} c_X X |\Delta\rangle. \quad (21)$$

We call the operators $X \in \mathcal{C}_l^p$ simply the *terms* of Ψ_l^p and the coefficients c_X its *coefficients*. Terms X with non-trivial coefficients c_X are called *non-trivial terms* of Ψ_l^p .

For each of the two types of twisted Verma modules $\mathcal{V}_\Delta$ and $\mathcal{V}_{\frac{c}{24}}^G$ we can thus think of two types of singular vectors $\Psi_l^\pm$ and $\Psi_{\frac{c}{24}-\Delta}^{\pm G}$ (both). However, the latter does not exist in G -closed Verma modules, as G -closed Verma modules $\mathcal{V}_{\frac{c}{24}}$ contain at level 0 only the highest weight vector^h.

To conclude this section, we should mention that in the case of a G -closed Verma module $\mathcal{V}_{\frac{c}{24}}^G$ the basis Eq. (20) simplifies to

$$\mathcal{B}_{\frac{c}{24}}^G = \left\{ X \left| \frac{c}{24} \right\rangle : X \in \mathcal{C}_l^{\pm G}, l \in \mathbb{N}_0^{1/2}, \right\}, \quad (22)$$

where

$$\begin{aligned} \mathcal{C}_l^{\pm G} = & \left\{ S_{k,p} T_{-\frac{1}{2}}^{2l-2k-r_1} G_{-\frac{1}{2}}^{r_1} : S_{k,p} \in \mathcal{S}_k^p, k \in \mathbb{N}_0^{1/2}, r_1 \in \{0, 1\}, \right. \\ & \left. 2l - 2k - r_1 \geq 0, p(-1)^{r_1} = \pm 1 \right\}, l \in \mathbb{N}_0^{1/2}. \end{aligned} \quad (23)$$

All other definitions and results carry over to the G -closed case simply by considering $\Delta = \frac{c}{24}$.

3 Adapted orderings and singular dimensions

A key question for the theory of highest weight representations is the dimension of any space of singular vectors that may appear at a given level $l \in \mathbb{N}_0^{1/2}$ with fixed parity $p \in \{\pm\}$ in a Verma module $\mathcal{V}_\Delta$. In Ref. 16 we have shown that these dimensions, called singular dimensions, are closely related to the ordering kernel of an adapted ordering on the set $\mathcal{C}_l^p$. In fact, it is the number of elements of the ordering kernel that sets an upper limit on the singular dimensions of a Verma module. Before we introduce an adapted ordering $\mathcal{O}$ on $\mathcal{C}_l^\pm$ and compute its ordering kernel as defined in Ref. 16, we shall first review a few results on adapted orderings that are crucial for our reasoning. An adapted ordering is simply a one-sided bounded total ordering on $\mathcal{C}_l^p$ satisfying the requirements specified in the following definition.

Definition 3.A *A total ordering $\mathcal{O}$ on $\mathcal{C}_l^p$ with global minimum is called adapted to the subset $\mathcal{C}_l^{pA} \subset \mathcal{C}_l^p$, in the Verma module $\mathcal{V}_\Delta$ with annihilation operators $\mathcal{K} = \mathcal{J}_2^+$, if for any element $X_0 \in \mathcal{C}_l^{pA}$ at least one annihilation operator $\Gamma \in \mathcal{K}$ exists for which*

$$\Gamma X_0 |\Delta\rangle = \sum_{X \in \mathcal{B}_\Delta} c_X^{\Gamma X_0} X \quad (24)$$

contains a non-trivial term $\tilde{X} \in \mathcal{B}_\Delta$ (i.e. $c_{\tilde{X}}^{\Gamma X_0} \neq 0$) such that for all $Y \in \mathcal{C}_l^p$ with $X_0 <_{\mathcal{O}} Y$ and $X_0 \neq Y$ the coefficient $c_{\tilde{X}}^{\Gamma Y}$ in

$$\Gamma Y |\Delta\rangle = \sum_{X \in \mathcal{B}_\Delta} c_X^{\Gamma Y} X \quad (25)$$

is trivial: $c_{\tilde{X}}^{\Gamma Y} = 0$. The complement of $\mathcal{C}_l^{pA}$, $\mathcal{C}_l^{pK} = \mathcal{C}_l^p \setminus \mathcal{C}_l^{pA}$ is the kernel with respect to the ordering $\mathcal{O}$ in the Verma module $\mathcal{V}_\Delta$.

^hThis resembles very much the result, for the Neveu-Schwarz and the topological $N=2$ algebras, that chiral singular vectors do not exist in chiral Verma modules.

If we replace in definition 3.A $\mathcal{C}_l^p$ by $\mathcal{C}_l^{pG}$, $\mathcal{C}_l^{pA}$ by $\mathcal{C}_l^{pGA}$, $\mathcal{V}_\Delta$ by $\mathcal{V}_{\frac{c}{24}}^G$, and $\mathcal{B}_{\frac{c}{24}}$ by $\mathcal{B}_{\frac{c}{24}}^G$, then we obtain the corresponding definition for a G -closed adapted ordering $\mathcal{O}_G$ with ordering kernel $\mathcal{C}_l^{pGK}$. For both cases, $\mathcal{O}$ and $\mathcal{O}_G$, if we replace the annihilation conditions by $\mathcal{K} = \mathcal{J}_2^+ \oplus \{G_0\}$ we obtain the corresponding definition for G -closed vectors at level l , parity p .

Our aim is to introduce an adapted ordering on $\mathcal{C}_l^p$ such that the resulting ordering kernels are as small as possible. There are two main reasons to proceed in this way. First, for a given singular vector Ψ_l^p of $\mathcal{V}_\Delta$ the coefficients of the terms corresponding to the ordering kernel $\mathcal{C}_l^{pK}$ determine the singular vector uniquely, i.e. if the normal forms of two singular vectors agree on the ordering kernel then the singular vectors are identical. This also implies that for a given singular vector Ψ_l^p at least one element of the adapted kernel must be a non-trivial term of Ψ_l^p , otherwise Ψ_l^p would agree on the whole adapted kernel with the trivial vector and would therefore be trivial itself. Secondly, the number of elements in the ordering kernel $\mathcal{C}_l^{pK}$ is an upper limit for the singular dimensions of $\mathcal{V}_\Delta$ at level l , parity p . These results are the content of two theorems which we proved in a very general setting in Ref. 16. Due to their importance for our later considerations we repeat these theorems here adapted to the particular case of the twisted $N = 2$ algebra $\mathcal{J}_2$.

Theorem 3.B *Let $\mathcal{O}$ denote an adapted ordering on $\mathcal{C}_l^{pA}$ at level l , parity p , with ordering kernel $\mathcal{C}_l^{pK}$ for a given Verma module $\mathcal{V}_\Delta$ and annihilation operators $\mathcal{K} = \mathcal{J}_2^+$. If the normal form of two vectors $\Psi_l^{p,1}$ and $\Psi_l^{p,2}$ at the same level l and parity p , both satisfying the highest weight conditions, have $c_X^1 = c_X^2$ for all terms $X \in \mathcal{C}_l^{pK}$, then*

$$\Psi_l^{p,1} \equiv \Psi_l^{p,2}. \quad (26)$$

Theorem 3.C *Let $\mathcal{O}$ denote an adapted ordering on $\mathcal{C}_l^{pA}$ at level l , parity p , with ordering kernel $\mathcal{C}_l^{pK}$ for a given Verma module $\mathcal{V}_\Delta$ and annihilation operators $\mathcal{K} = \mathcal{J}_2^+$. If the ordering kernel $\mathcal{C}_l^{pK}$ has n elements, then there are at most n linearly independent singular vectors Ψ_l^p in $\mathcal{V}_\Delta$ at level l with parity p .*

Again, we can replace $\mathcal{O}$ on $\mathcal{C}_l^p$ by $\mathcal{O}_G$ on $\mathcal{C}_l^{pG}$ in order to obtain both theorems for G -closed Verma modules $\mathcal{V}_{\frac{c}{24}}^G$. If we extend the annihilation operators from $\mathcal{K} = \mathcal{J}_2^+$ to $\mathcal{K} = \mathcal{J}_2^+ \oplus \{G_0\}$, which normally leads to a smaller ordering kernel, then both theorems hold with this smaller ordering kernel for G -closed singular vectors Ψ_l^{pG} .

4 Adapted ordering for the twisted $N = 2$ algebra

In this section we will define an adapted ordering on $\mathcal{C}_l^p$, given by Eq. (18), and compute its ordering kernel which will later turn out to be the smallest possible ordering kernel. We will conclude this section by transferring these results to the G -closed case $\mathcal{C}_l^{pG}$. For convenience, we shall first give an ordering on the sets $\mathcal{L}_m$, $\mathcal{T}_s$, and $\mathcal{G}_k$.

Definition 4.A *Let Y denote either $\mathcal{L}$, $\mathcal{T}$, or $\mathcal{G}$, (but the same throughout this definition) and take two elements Y_i , $i = 1, 2$, such that $Y_i = Z_{-k_{\parallel Y_i}}^i \dots Z_{-k_1}^i$, $|Y_i|_L = k^i$ for $k^i \in \mathbb{N}_0^{1/2}$, or $Y_i = 1$, $i = 1, 2$, with $Z_{-k_j}^i$ being generators of the type $L_{-k_j^i}$, $T_{-k_j^i}$, or $G_{-k_j^i}$ depending on whether Y denotes $\mathcal{L}$, $\mathcal{T}$, or $\mathcal{G}$ respectively. For $Y_1 \neq Y_2$ we compute the indexⁱ $j_0 = \min\{j :$*

ⁱFor subsets of $\mathbb{N}$ we define $\min \emptyset = 0$.

$k_j^1 - k_j^2 \neq 0, j = 1, \dots, \min(\|Y_1\|, \|Y_2\|)\}$. If non-trivial, j_0 is the index for which the levels of the generators in Y_1 and Y_2 first disagree when read from the right to the left. For $j_0 > 0$ we then define

$$Y_1 <_Y Y_2 \quad \text{if} \quad k_{j_0}^1 < k_{j_0}^2. \quad (27)$$

If, however, $j_0 = 0$, we set

$$Y_1 <_Y Y_2 \quad \text{if} \quad \|Y_1\| > \|Y_2\|. \quad (28)$$

In order to get more familiar with these definitions let us consider the following examples:

$$\begin{aligned} L_{-2}L_{-2}L_{-1} &<_L L_{-5}L_{-1} \quad , \quad \text{where } j_0 = 2 \quad , \\ G_{-3}G_{-\frac{5}{2}}G_{-1} &<_G G_{-\frac{5}{2}}G_{-1} \quad , \quad \text{where } j_0 = 0 \quad , \\ T_{-\frac{3}{2}} &<_T 1 \quad , \quad \text{where } j_0 = 0 \quad . \end{aligned} \quad (29)$$

We can now define an ordering on $\mathcal{C}_l^p$ which will turn out to be adapted with a sufficiently small ordering kernel.

Definition 4.B On the set $\mathcal{C}_l^p$, $l \in \mathbb{N}^{1/2}$, $p \in \{\pm\}$ we introduce the total ordering $\mathcal{O}$. For two elements $X_1, X_2 \in \mathcal{C}_l^p$, $X_1 \neq X_2$ with $X_i = L^i T^i G^i T_{-\frac{1}{2}}^{n_i} G_{-\frac{1}{2}}^{r_i} G_0^{r_i}$, $L^i \in \mathbb{L}_{m_i}$, $T^i \in \mathbb{T}_{s_i}$, $G^i \in \mathbb{G}_{k_i}$ for some $m_i, n_i \in \mathbb{N}_0$, $s_i, k_i \in \mathbb{N}^{1/2}$, $r_1^i, r_2^i \in \{0, 1\}$, $i = 1, 2$, we define

$$X_1 <_{\mathcal{O}} X_2 \quad \text{if} \quad n^1 > n^2. \quad (30)$$

For $n^1 = n^2$ we set

$$X_1 <_{\mathcal{O}} X_2 \quad \text{if} \quad r_1^1 > r_1^2. \quad (31)$$

If $r_1^1 = r_1^2$ then we set

$$X_1 <_{\mathcal{O}} X_2 \quad \text{if} \quad G^1 <_G G^2. \quad (32)$$

In the case that $G^1 = G^2$ we then define

$$X_1 <_{\mathcal{O}} X_2 \quad \text{if} \quad L^1 <_L L^2. \quad (33)$$

If further $L^1 = L^2$ we set

$$X_1 <_{\mathcal{O}} X_2 \quad \text{if} \quad T^1 <_T T^2, \quad (34)$$

which finally has to give an answer. For $X_1 = X_2$ we define $X_1 <_{\mathcal{O}} X_2$ and $X_2 <_{\mathcal{O}} X_1$.

Definition 4.B is well-defined since one obtains an answer for any pair $X_1, X_2 \in \mathcal{C}_l^p$, $X_1 \neq X_2$ after going through EQS. (30)-(34), and hence the ordering $\mathcal{O}$ proves to be a total ordering on $\mathcal{C}_l^p$. Namely, if EQS. (30)-(34) do not give an answer on the ordering of X_1 and X_2 , then obviously X_1 and X_2 are of the form $X_i = L T G T_{-\frac{1}{2}}^{n_i} G_{-\frac{1}{2}}^{r_i} G_0^{r_i}$, with common L , T , G , n , and also r_1 . The fact that both X_1 and X_2 have the same parity p implies further $r_2^1 = r_2^2$ and ultimately $X_1 = X_2$.

It is easy to see that the $\mathcal{O}$ -smallest element of $\mathcal{C}_l^+$ is $T_{-\frac{1}{2}}^{2l}$ followed by $T_{-\frac{1}{2}}^{2l-1} G_{-\frac{1}{2}} G_0$ whilst the $\mathcal{O}$ -smallest element of $\mathcal{C}_l^-$ is $T_{-\frac{1}{2}}^{2l} G_0$ followed by $T_{-\frac{1}{2}}^{2l-1} G_{-\frac{1}{2}}$. We will now show that the

ordering $\mathcal{O}$ is adapted and we will compute the ordering kernels. It will turn out that the elements of the ordering kernels are exactly the four elements just mentioned and therefore they identify all singular vectors according to theorem 3.B. In section 6 we shall give examples of singular vectors at low levels which show that these are indeed the smallest ordering kernels for general values of Δ .

Theorem 4.C *For all central terms $c \in \mathbb{C}$, the ordering $\mathcal{O}$ is adapted to $\mathcal{C}_l^p$ for all Verma modules $\mathcal{V}_\Delta$, for all levels l with $l \in \mathbb{N}_0^{1/2}$, and for both parities $p \in \{\pm\}$. The ordering kernels are given by the following tables for all levels^j l .*

$(-1)^F$	ordering kernel
+1	$\{T_{-\frac{1}{2}}^{2l}, T_{-\frac{1}{2}}^{2l-1} G_{-\frac{1}{2}} G_0\}$
-1	$\{T_{-\frac{1}{2}}^{2l} G_0, T_{-\frac{1}{2}}^{2l-1} G_{-\frac{1}{2}}\}$

TAB. a Ordering kernels for $\mathcal{O}$, annihilation operators J_2^+ .

$(-1)^F$	ordering kernel
+1	$\{T_{-\frac{1}{2}}^{2l}\}$
-1	$\{T_{-\frac{1}{2}}^{2l} G_0\}$

TAB. b Ordering kernels for $\mathcal{O}$, annihilation operators J_2^+ and G_0 .

For the proof of theorem 4.C we follow the lines of the proof for the topological $N = 2$ algebra given in Ref. 16. There, however, our strategy for the proof was to find annihilation operators that were able to produce an additional operator L_{-1} which would create a term that cannot be obtained by any $\mathcal{O}$ -bigger term. Obviously, in our present case $T_{-\frac{1}{2}}$ has taken over the rôle that was played by L_{-1} in the topological $N = 2$ case. (As a matter of fact this is the first time that this rôle is played by a generator different from L_{-1}). The following proof makes clear that in this case the adapted ordering method works for powers of $T_{-\frac{1}{2}}$.

Proof: Let us start with the most general term X_0 at level l , parity p , in $\mathcal{C}_l^p$, $l \in \mathbb{N}_0^{1/2}$, $p \in \{\pm\}$.

$$X_0 = L^0 T^0 G^0 T_{-\frac{1}{2}}^{r_1} G_{-\frac{1}{2}}^{r_2} G_0^{r_2} \in \mathcal{C}_l^p, \quad (35)$$

with

$$\begin{aligned} L^0 &= L_{-m_{\parallel L^0}} \dots L_{-m_1} \in \mathbb{L}_m, \\ T^0 &= T_{-s_{\parallel T^0}} \dots T_{-s_1} \in \mathbb{T}_s, \\ G^0 &= G_{-k_{\parallel G^0}} \dots G_{-k_1} \in \mathbb{G}_k, \end{aligned} \quad (36)$$

$m, n \in \mathbb{N}_0$, $s, k \in \mathbb{N}_0^{1/2}$, $r_1, r_2 \in \{0, 1\}$, such that $l = m + s + k + \frac{n}{2} + \frac{r_1}{2}$ and $p = (-1)^{\|G^0\| + r_1 + r_2}$. We then construct the vector $\Psi^0 = X_0 | \Delta \rangle$.

In the case $G^0 \neq 1$ we look at G_{-k_1} which necessarily has $k_1 > \frac{1}{2}$ and consider the positive operator $G_{k_1 - \frac{1}{2}}$. The commutation relations of G_{-k_1} and $G_{k_1 - \frac{1}{2}}$ create a generator $T_{-\frac{1}{2}}$ (note

^jNote that for level $l = 0$ some of the kernel elements obviously do not exist.

that $2k_1 - \frac{1}{2} > 0$). Therefore, acting with the positive operator $G_{k_1 - \frac{1}{2}}$ on $\Psi_0 = X_0 |\Delta\rangle$ we obtain a non-trivial term X^G in $G_{k_1 - \frac{1}{2}} \Psi^0$ of the form

$$\begin{aligned} X^G &= L^0 T^0 \tilde{G}^0 T_{-\frac{1}{2}}^{n+1} G_{-\frac{1}{2}}^{r_1} G_0^{r_2} \in \mathcal{C}_{l-k_1+\frac{1}{2}}^{-p}, \\ \tilde{G}^0 &= G_{-k_{\parallel G^0}} \dots G_{-k_2}, \end{aligned} \quad (37)$$

or $\tilde{G}^0 = 1$ in the case $\|G^0\| = 1$. Any other term $Y \in \mathcal{C}_l^p$ which also produces X^G under the action of $G_{k_1 - \frac{1}{2}}$ also creates at least one generator $T_{-\frac{1}{2}}$ under the action of $G_{k_1 - \frac{1}{2}}$ or is already $\mathcal{O}$ -smaller than X_0 since Y would already have more generators $T_{-\frac{1}{2}}$ than X_0 . The latter case is irrelevant for us in view of definition 3.A on adapted orderings. The action of one positive operator can however create at most one new generator. We can therefore restrict ourselves to terms Y of the form

$$Y = L^Y T^Y G^Y T_{-\frac{1}{2}}^n G_{-\frac{1}{2}}^{r_1^Y} G_0^{r_2^Y} \in \mathcal{C}_l^p, \quad (38)$$

with the same number n of generators $T_{-\frac{1}{2}}$ than X_0 . If $G_{k_1 - \frac{1}{2}}$ creates X^G acting on Y , then the additional $T_{-\frac{1}{2}}$ could only be created in three ways. First, $G_{k_1 - \frac{1}{2}}$ commutes with generators in G^Y and creates $T_{-\frac{1}{2}}$. Second, $G_{k_1 - \frac{1}{2}}$ commutes with generators in $L^Y T^Y$ and creates $G_{k'}$ with $k' < k_1$ which furthermore commutes with generators in G^Y to create $T_{-\frac{1}{2}}$. Third, $G_{k_1 - \frac{1}{2}}$ commutes with generators in $L^Y T^Y G^Y$ and creates G_0 which then commutes with $G_{-\frac{1}{2}}$ in the case of $r_1^Y = 1$ to create $T_{-\frac{1}{2}}$. In the third case $r_1^Y = 1$ would obviously lead to Y being $\mathcal{O}$ -smaller than X_0 because then $r_1 = 0$ (as X^G is the term produced from Y in this way), and due to Eq. (31). In the first and second cases we obviously need to have a generator G_{-k} in G^Y with $\frac{1}{2} < k < k_1$, or G_{-k_1} itself is contained in G^Y and $G_{k_1 - \frac{1}{2}}$ commutes only with this generator. In the latter case X_0 would consequently equal Y and in the former case Y is again $\mathcal{O}$ -smaller than X_0 according to Eq. (32). Hence, we have shown that the ordering $\mathcal{O}$ is adapted on the set of terms X_0 of the form Eq. (35) with $G^0 \neq 1$.

Let us therefore continue with terms X_0 of the form

$$X_0 = L^0 T^0 T_{-\frac{1}{2}}^n G_{-\frac{1}{2}}^{r_1} G_0^{r_2} \in \mathcal{C}_l^p. \quad (39)$$

If $L^0 \neq 1$ we consider the positive operator $T_{m_1 - \frac{1}{2}}$. Acting with $T_{m_1 - \frac{1}{2}}$ on $\Psi_0 = X_0 |\Delta\rangle$ creates a non-trivial term

$$\begin{aligned} X^T &= \tilde{L}^0 T^0 T_{-\frac{1}{2}}^{n+1} G_{-\frac{1}{2}}^{r_1} G_0^{r_2} \in \mathcal{C}_{l-m_1+\frac{1}{2}}^p, \\ \tilde{L}^0 &= L_{-m_{\parallel L^0}} \dots L_{-m_2}, \end{aligned} \quad (40)$$

or $\tilde{L} = 1$ in the case $\|L^0\| = 1$. Again, we find that any term $Y \in \mathcal{C}_l^p$ which is not $\mathcal{O}$ -smaller than X_0 but still produces X^T under the action of $T_{m_1 - \frac{1}{2}}$ needs to create exactly one generator $T_{-\frac{1}{2}}$ and thus $n^Y = n$. As in the case before we find $r_1^Y = r_1$, and also $G^Y = 1$, otherwise $Y <_{\mathcal{O}} X_0$. We can hence concentrate on terms Y of the form

$$Y = L^Y T^Y T_{-\frac{1}{2}}^n G_{-\frac{1}{2}}^{r_1} G_0^{r_2} \in \mathcal{C}_l^p, \quad (41)$$

where also $r_2^Y = r_2$ due to parity equality with X_0 . The only way of creating $T_{-\frac{1}{2}}$ from Y under the action of $T_{m_1 - \frac{1}{2}}$ is via commutation with generators in L^Y as the commutation with

generators in T^Y does not produce any new generators. Thus, $T_{m_1-\frac{1}{2}}$ can produce $T_{-\frac{1}{2}}$ in one step by commuting with L_{-m_1} if this is contained in L^Y . In this case, however, Y turns out to be equal to X_0 . Another possibility is that $T_{m_1-\frac{1}{2}}$ produces $T_{-\frac{1}{2}}$ in more than one step by commuting with generators in L^Y which obviously requires L^Y to contain generators L_{-m} with $1 \leq m < m_1$ and therefore $Y <_{\mathcal{O}} X_0$ due to EQ. (33). We therefore have shown that the ordering $\mathcal{O}$ is adapted on the set of terms X_0 of the form EQ. (35) with $G^0 \neq 1$ or $L^0 \neq 1$.

For the next step we can thus assume that X_0 is of the form

$$X_0 = T^0 T_{-\frac{1}{2}}^n G_{-\frac{1}{2}}^{r_1} G_0^{r_2} \in \mathcal{C}_l^p . \quad (42)$$

If $T^0 \neq 1$ we take the positive operator $L_{s_1-\frac{1}{2}}$. Acting with $L_{s_1-\frac{1}{2}}$ on $\Psi_0 = X_0 |\Delta\rangle$ creates a non-trivial term

$$\begin{aligned} X^L &= \tilde{T}^0 T_{-\frac{1}{2}}^{n+1} G_{-\frac{1}{2}}^{r_1} G_0^{r_2} \in \mathcal{C}_{l-s_1+\frac{1}{2}}^p , \\ \tilde{T}^0 &= T_{-s_{\|T^0\|}} \dots T_{-s_2} , \end{aligned} \quad (43)$$

or $\tilde{T} = 1$ in the case $\|T^0\| = 1$. Similar arguments as above allow us to restrict ourselves to terms Y of the form

$$Y = T^Y T_{-\frac{1}{2}}^n G_{-\frac{1}{2}}^{r_1} G_0^{r_2} \in \mathcal{C}_l^p , \quad (44)$$

as all other terms Y that also produce X^L under the action of $L_{s_1-\frac{1}{2}}$ would automatically be $\mathcal{O}$ -smaller than X_0 . Similarly as before, the only way of creating $T_{-\frac{1}{2}}$ from Y under the action of $L_{s_1-\frac{1}{2}}$ is by commuting $L_{s_1-\frac{1}{2}}$ with generators in T^Y . Exactly the same arguments as for L^Y together with EQ. (34) show that either $Y = X_0$ or $Y <_{\mathcal{O}} X_0$. Hence, the ordering $\mathcal{O}$ is adapted on the set of terms X_0 of the form EQ. (35) with $G^0 \neq 1$ or $L^0 \neq 1$ or $T^0 \neq 1$. This already proves the ordering kernels of table TAB. a.

Finally, if we also have G_0 as annihilation operator, we can then act with G_0 on terms X_0 of the form

$$X_0 = T_{-\frac{1}{2}}^n G_{-\frac{1}{2}} G_0^{r_2} \in \mathcal{C}_l^p , \quad (45)$$

which creates a non-trivial term

$$X' = T_{-\frac{1}{2}}^{n+1} G_0^{r_2} \in \mathcal{C}_l^{-p} . \quad (46)$$

Obviously, the only way of creating $T_{-\frac{1}{2}}$ by commuting G_0 with products of generators is by commuting it with $G_{-\frac{1}{2}}$. Therefore any other term Y that creates X' under the action of G_0 and is not $\mathcal{O}$ -smaller than X_0 would also need to create one $T_{-\frac{1}{2}}$ and thus also have $r_1^Y = r_1 = 1$, besides $n^Y = n$. Trivially, we find in this case that $Y = X_0$. This proves the ordering kernel of table TAB. b which finally completes our proof of theorem 4.C. $\square$

To conclude this subsection we mention that we can easily transfer these results to G -closed Verma modules. It is fairly straightforward to see that the above proof also holds if we set everywhere $r_2 \equiv 0$. This however leads exactly to the G -closed case. Therefore, the same ordering $\mathcal{O}$ of definition 4.B will also serve as ordering $\mathcal{O}_G$ on $\mathcal{C}_l^{pG}$ in the case of a G -closed Verma module $\mathcal{V}_{\frac{c}{24}}^G$. Setting $r_2 \equiv 0$ everywhere in the proof of theorem 4.C obviously reduces the ordering kernels. One easily finds the following theorem.

Theorem 4.D For all central terms $c \in \mathbb{C}$, the ordering $\mathcal{O}$ is adapted to $\mathcal{C}_l^{pG}$ for all Verma modules $\mathcal{V}_{\frac{c}{24}}^G$, for all levels l with $l \in \mathbb{N}_0^{1/2}$, and for both parities $p \in \{\pm\}$. Ordering kernels are given by the following tables for all levels l .

$(-1)^F$	ordering kernel
+1	$\{T_{-\frac{1}{2}}^{2l}\}$
-1	$\{T_{-\frac{1}{2}}^{2l-1}G_{-\frac{1}{2}}\}$

TAB. c Ordering kernels for $\mathcal{O}$ on $\mathcal{C}_l^{pG}$, annihilation operators J_2^+ .

5 Singular dimensions and singular vectors of the twisted $N = 2$ algebra

As a consequence of the results of the previous section we will now be able to give an upper limit for the dimensions of a singular vector space at a given level $l \in \mathbb{N}_0^{1/2}$ with fixed parity $p \in \{\pm\}$. In addition, we will use the ordering kernels in order to uniquely identify all singular vectors (primitive as well as secondary) of the twisted $N = 2$ superconformal algebra. In section 6 we shall give explicit examples which prove that for particular Verma modules the upper limits are indeed the maximal singular dimensions.

Theorem 3.C tells us that an upper limit for the dimension of a space of singular vectors at fixed level l and parity p is given by the number of elements in the corresponding ordering kernel $\mathcal{C}_l^{pK}$. Using the tables of ordering kernels of theorem 4.C and theorem 4.D we can easily give the following maximal singular dimensions.

Theorem 5.A For singular vectors Ψ_l^p and G -closed singular vectors Ψ_l^{pG} in the Verma modules $\mathcal{V}_\Delta$ and $\mathcal{V}_{\frac{c}{24}}^G$ (both), one finds the following upper limits for the number of linearly independent singular vectors at the same level $l \in \mathbb{N}_0^{1/2}$ and with the same parity $p \in \{\pm\}$.

	$p = +1$	$p = -1$
$\Psi_l^p \in \mathcal{V}_\Delta$	2	2
$\Psi_l^p \in \mathcal{V}_{\frac{c}{24}}^G$	1	1
$\Psi_{\frac{c}{24}-\Delta}^{pG} \in \mathcal{V}_\Delta$	1	1
$\Psi_0^{pG} \in \mathcal{V}_{\frac{c}{24}}^G$	0	0

TAB. d Maximal dimensions for singular spaces of the twisted $N = 2$ superconformal algebra.

Surely, not at each level the singular spaces will have the dimensions given in the previous tables. In fact, for most levels in most Verma modules there will not be any singular vectors at all. The determinant formulae of the inner product matrix⁸, given in section 2, indicate in which Verma modules and at which levels singular vectors appear (for the first time). In the following section we shall give examples of these singular vectors for low levels. From theorem 3.B we know that singular vectors can only exist if they contain in their normal form at least one non-trivial term of the corresponding ordering kernel of theorem 4.C or 4.D. Furthermore, if the normal forms of two singular vectors agree in the coefficients of the terms of the ordering kernel, then the singular vectors are identical. This allows us to identify existing singular vectors simply by their coefficients with respect to the terms of the ordering kernel. For the cases of TAB. d with singular dimension 1 this implies that two singular vectors are already proven to

be proportional if we can simply show that they are on the same level with the same parity. In addition, if the coefficient of the term of the ordering kernel vanishes, then the singular vector is trivial. Thus, the coefficient of $T_{-\frac{1}{2}}^{2l}$ is already sufficient in order to see if a singular vector is trivial, in the case of $p = +$ (see tables TAB. b and TAB. c), whereas the coefficient of $T_{-\frac{1}{2}}^{2l}G_0$ or $T_{-\frac{1}{2}}^{2l-1}G_{-\frac{1}{2}}$ is sufficient in order to decide if a singular vector is trivial in the case $p = -$. When the ordering kernel is two-dimensional we need the coefficients of the two terms in order to uniquely define a singular vector. This justifies to introduce an appropriate notation, as follows.

Definition 5.B Let $\Psi_l^\pm = \theta_l^\pm |\Delta\rangle$ be a singular vector in $\mathcal{V}_\Delta$ at level $l \in \mathbb{N}_0^{1/2}$ with parity $\{\pm\}$. The normal form of Ψ_l^+ is completely determined by the coefficients of $T_{-\frac{1}{2}}^{2l}$ and $T_{-\frac{1}{2}}^{2l-1}G_{-\frac{1}{2}}G_0$, whilst the normal form of Ψ_l^- is determined by the coefficients of $T_{-\frac{1}{2}}^{2l}G_0$ and $T_{-\frac{1}{2}}^{2l-1}G_{-\frac{1}{2}}$. We thus use the following notation

$$(a, b)_l^+ = \theta_l^+ = aT_{-\frac{1}{2}}^{2l} + bT_{-\frac{1}{2}}^{2l-1}G_{-\frac{1}{2}}G_0 + \dots, \quad (47)$$

$$(a, b)_l^- = \theta_l^- = aT_{-\frac{1}{2}}^{2l}G_0 + bT_{-\frac{1}{2}}^{2l-1}G_{-\frac{1}{2}} + \dots, \quad (48)$$

for the singular vector operators $\theta_l^\pm$ given in their normal form, where $a, b \in \mathbb{C}$.

Notice that definition 5.B only makes sense if $\Psi_l^\pm$ is a singular vector. Certainly for most pairs $a, b \in \mathbb{C}$ singular vectors do not exist.

The advantage of this notation lies in the fact that we can easily obtain multiplication rules for singular vector operators and construct descendant singular vectors. (In the context of embedding diagrams, this issue will be analysed in more detail in a forthcoming publication). For example, the product of the first terms of the singular vector operators

$$\begin{aligned} (a_1T_{-\frac{1}{2}}^{2l_1} + b_1T_{-\frac{1}{2}}^{2l_1-1}G_{-\frac{1}{2}}G_0 + \dots)(a_2T_{-\frac{1}{2}}^{2l_2} + b_2T_{-\frac{1}{2}}^{2l_2-1}G_{-\frac{1}{2}}G_0 + \dots) = \\ a_1a_2T_{-\frac{1}{2}}^{2(l_1+l_2)} + (a_1b_2 + a_2b_1 - \frac{b_1b_2}{2})T_{-\frac{1}{2}}^{2(l_1+l_2)-1}G_{-\frac{1}{2}}G_0 + \dots, \end{aligned} \quad (49)$$

gives us the multiplication rule for $(a_1, b_1)_{l_1}^+(a_2, b_2)_{l_2}^+$ provided the conformal weights match: $\Delta_2 + l_2 = \Delta_1$. In the same way we can easily find similar rules for products involving negative parity operators. The following theorem summarizes these results.

Theorem 5.C Given two singular vector operators $\theta_{l_1}^{p_1}$ and $\theta_{l_2}^{p_2}$ for the Verma modules $\mathcal{V}_{\Delta_1}$ and $\mathcal{V}_{\Delta_2}$ with $\Delta_2 + l_2 = \Delta_1$ then $\theta_{l_1}^{p_1}\theta_{l_2}^{p_2}|\Delta_2\rangle$ is either trivial or singular in $\mathcal{V}_{\Delta_2}$ at level $l_1 + l_2$ with parity p_1p_2 . The resulting singular vector can be expressed in the following way, depending on the multiplication rules which in turn depend on the parities:

$$(a_1, b_1)_{l_1}^+(a_2, b_2)_{l_2}^+|\Delta_2\rangle = (a_1a_2, a_1b_2 + a_2b_1 - \frac{b_1b_2}{2})_{l_1+l_2}^+|\Delta_2\rangle, \quad (50)$$

$$(a_1, b_1)_{l_1}^+(a_2, b_2)_{l_2}^-|\Delta_2\rangle = (a_1a_2, a_1b_2 + a_2b_1[\Delta_2 - \frac{c}{24}] - \frac{b_1b_2}{2})_{l_1+l_2}^-|\Delta_2\rangle, \quad (51)$$

$$(a_1, b_1)_{l_1}^-(a_2, b_2)_{l_2}^+|\Delta_2\rangle = (a_1a_2 - \frac{a_1b_2}{2}, -2l_2a_1b_2 + a_2b_1 - a_1b_2[\Delta_2 - \frac{c}{24}])_{l_1+l_2}^-|\Delta_2\rangle, \quad (52)$$

$$(a_1, b_1)_{l_1}^-(a_2, b_2)_{l_2}^-|\Delta_2\rangle = (a_1a_2[\Delta_2 - \frac{c}{24}] - \frac{a_1b_2}{2}, a_2b_1 - 2l_2a_1b_2 - a_1b_2)_{l_1+l_2}^+|\Delta_2\rangle. \quad (53)$$

For the vectors $\Psi_{G,l}^\pm$ and ${}_G\Psi_l^\pm$, introduced in Eqs. (8)-(11), we can derive the corresponding expressions by looking at operator products of the type

$$(aT_{-\frac{1}{2}}^{2l} + bT_{-\frac{1}{2}}^{2l-1}G_{-\frac{1}{2}}G_0 + \dots)G_0 = aT_{-\frac{1}{2}}^{2l}G_0 + bT_{-\frac{1}{2}}^{2l-1}G_{-\frac{1}{2}}(L_0 - \frac{c}{24}) + \dots, \quad (54)$$

and similar products for negative parity vectors. We obtain the following results for $\Psi_{G,l}^\pm = (a, b)_{G,l}^\pm |\Delta\rangle$ and ${}_G\Psi_l^\pm = {}_G(a, b)_l^\pm |\Delta\rangle$:

$${}_G(a, b)_l^- |\Delta\rangle = (a - \frac{b}{2}, -b[\Delta - \frac{c}{24}] - 2la)_l^- |\Delta\rangle, \quad (55)$$

$$(a, b)_{G,l}^- |\Delta\rangle = (a, b[\Delta - \frac{c}{24}])_l^- |\Delta\rangle, \quad (56)$$

$${}_G(a, b)_l^+ |\Delta\rangle = (a[\Delta - \frac{c}{24}] - \frac{b}{2}, -2la - b)_l^+ |\Delta\rangle, \quad (57)$$

$$(a, b)_{G,l}^+ |\Delta\rangle = (a[\Delta - \frac{c}{24}], b)_l^+ |\Delta\rangle. \quad (58)$$

6 Level $1/2$, level 1, and level $3/2$ singular vectors.

Before we proceed in analysing the structure of the singular vectors we shall first give all singular vectors at the lower levels $1/2$, 1 and $3/2$. These examples already prove that the upper limits given for the singular dimensions are in fact not only upper limits but actually maximal singular dimensions for certain Verma modules. The determinant formulae⁸, given in section 2, show the existence of the parity – singular vector Ψ_0^- at level 0 in the Verma modules $\mathcal{V}_{\frac{c}{24}}$, and also show the existence of the singular vectors $\Psi_{r,s}^\pm$ at level $\frac{rs}{2}$ with parity $\pm$, $r, s \in \mathbb{N}$, s odd, in the Verma modules $\mathcal{V}_{\Delta_{r,s}(t)}$ with:

$$\Delta_{r,s}(t) = -\frac{(rt+s)^2}{8t} + \frac{t}{8} + \frac{1}{8}, \quad (59)$$

$$c = 3 + 3t. \quad (60)$$

These singular vectors $\Psi_{r,s}^\pm$ are generically primitive. However in the case of intersections, with two or more of them in the same Verma module, some of these singular vectors are actually secondary.

By explicit computer calculations²⁹ one finds the following singular vectors (one-parameter families of singular vectors, to be precise) for parity $+$. The corresponding results for parity $-$ can easily be derived using the multiplication rules Eqs. (55)-(58). These singular vectors are given in appendix A.

Level $\frac{1}{2}$:

$$\Psi_{1,1}^+(t) = \left\{ (t+1)T_{-\frac{1}{2}} + 4tG_{-\frac{1}{2}}G_0 \right\} \left| -\frac{t+1}{8t} \right\rangle. \quad (61)$$

Level 1:

$$\begin{aligned} \Psi_{2,1}^+(t) = & \left\{ (2t+1)T_{-\frac{1}{2}}^2 + 8tT_{-\frac{1}{2}}G_{-\frac{1}{2}}G_0 + 2t(2t+1)L_{-1} \right. \\ & \left. - 4t(t+1)G_{-1}G_0 \right\} \left| -\frac{3t^2+3t+1}{8t} \right\rangle. \end{aligned} \quad (62)$$

Level $\frac{3}{2}$ has two different singular vectors:

$$\begin{aligned} \Psi_{3,1}^+(t) = & \left\{ (3t+1)T_{-\frac{1}{2}}^3 + 12tT_{-\frac{1}{2}}^2G_{-\frac{1}{2}}G_0 + 6t(3t+2)L_{-1}T_{-\frac{1}{2}} \right. \\ & - 4t(4t+3)G_{-1}T_{-\frac{1}{2}}G_0 + 8t^2L_{-1}G_{-\frac{1}{2}}G_0 + 2t(3t+1)G_{-1}G_{-\frac{1}{2}} \\ & \left. + 2t(t-1)(3t+1)T_{-\frac{3}{2}} + 8t(t^2-t-1)G_{-\frac{3}{2}}G_0 \right\} \left| -\frac{8t^2+5t+1}{8t} \right\rangle, \end{aligned} \quad (63)$$

$$\begin{aligned} \Psi_{1,3}^+(t) = & \left\{ (t+3)T_{-\frac{1}{2}}^3 + 4tT_{-\frac{1}{2}}^2G_{-\frac{1}{2}}G_0 - 2t(t+3)L_{-1}T_{-\frac{1}{2}} \right. \\ & - 4tG_{-1}T_{-\frac{1}{2}}G_0 - 8t^2L_{-1}G_{-\frac{1}{2}}G_0 + 2t(t+3)G_{-1}G_{-\frac{1}{2}} \\ & \left. - 4(t+3)T_{-\frac{3}{2}} - 8tG_{-\frac{3}{2}}G_0 \right\} \left| -\frac{5t+9}{8t} \right\rangle. \end{aligned} \quad (64)$$

We can use the notation introduced in definition 5.B of the previous section in order to identify the given examples of singular vectors in terms of their leading coefficients.

$$\Psi_{1,1}^+(t) = (t+1, 4t)_{\frac{1}{2}}^+ \left| -\frac{t+1}{8t} \right\rangle, \quad (65)$$

$$\Psi_{2,1}^+(t) = (2t+1, 8t)_1^+ \left| -\frac{3t^2+3t+1}{8t} \right\rangle, \quad (66)$$

$$\Psi_{3,1}^+(t) = (3t+1, 12t)_{\frac{3}{2}}^+ \left| -\frac{8t^2+5t+1}{8t} \right\rangle, \quad (67)$$

$$\Psi_{1,3}^+(t) = (t+3, 4t)_{\frac{3}{2}}^+ \left| -\frac{5t+9}{8t} \right\rangle. \quad (68)$$

We should note that at least at these levels there are no other singular vectors than the ones predicted by the determinant formula. In principle, the determinant formula only proves the existence of the singular vector at lowest level in a Verma module given by Eq. (59). Continuation arguments extend the existence to the cases where the curves of Eq. (59) intersect. The determinant formula does *a priori* not exclude the existence of so-called *isolated* singular vectors which are primitive singular vectors that may appear at higher levels and are always *hidden* behind a singular vector of type $\Psi_{r,s}^\pm$. As far as we know, such isolated singular vectors have never been found for any chiral algebra.

As these examples are the complete set of singular vectors at levels $\frac{1}{2}$, 1, or $\frac{3}{2}$ with positive parity, we observe that the singular spaces at level $\frac{1}{2}$ and level 1 are one-dimensional. However, at level $\frac{3}{2}$ we find that the conformal weights $\Delta_{3,1}(t)$ and $\Delta_{1,3}(t)$ intersect for $t = \pm 1$. In these cases the corresponding singular vectors are $\Psi_{3,1}^+(1) = (4, 12)_{\frac{3}{2}}^+ \left| -\frac{7}{4} \right\rangle$ and $\Psi_{1,3}^+(1) = (4, 4)_{\frac{3}{2}}^+ \left| -\frac{7}{4} \right\rangle$ for $t = 1$, and $\Psi_{3,1}^+(-1) = (-2, -12)_{\frac{3}{2}}^+ \left| \frac{1}{2} \right\rangle$ and $\Psi_{1,3}^+(-1) = (2, -4)_{\frac{3}{2}}^+ \left| \frac{1}{2} \right\rangle$ for $t = -1$. Obviously, for both values of t these singular vectors are linearly independent and thus span a two-dimensional singular vector space. Therefore we also find degenerate singular vectors for the twisted $N = 2$ superconformal algebra as already observed for the three isomorphic $N = 2$ superconformal algebras^{12, 20, 21}.

We should stress that in the Virasoro case for a given level $l \in \mathbb{N}_0$ we also have different one-parameter families of singular vectors $\xi_{p,q}(t)$ that intersect for certain values of t . However, for Virasoro Verma modules these singular vectors at such intersection points are always linearly

dependent. This follows indirectly from the proof of Feigin and Fuchs¹⁷ and directly from the results of Kent²⁶ (see also our results in Ref. 16). The same happens for the Verma modules of the $N = 1$ superconformal algebras. The degeneracy found for the Neveu-Schwarz $N = 2$ algebra¹² (and for the topological and Ramond $N = 2$ algebras^{16, 21}) is different in the sense that one single uncharged vector $\Psi_{r,s}^{NS}$ may on its own describe a two-dimensional space under certain conditions (+1 and -1 charged singular vectors intersecting in the same Verma module and the uncharged singular vector becoming a *secondary* singular vector of them). At these points $t = t_0$, $\Psi_{r,s}^{NS}(t_0)$ vanishes and the two-dimensional tangent space becomes a degenerate singular space. The degeneracy found for the twisted $N = 2$ algebra is thus of a different type since the singular vectors that intersect are generically *primitive*, like the vectors $\Psi_{3,1}^+(1)$ and $\Psi_{1,3}^+(1)$, and $\Psi_{3,1}^+(-1)$ and $\Psi_{1,3}^+(-1)$ given above. It is in fact unique of its kind in the sense that it has not been observed for any other chiral algebra so far in this form. In the following section we will conjecture general expressions for the twisted $N = 2$ singular vectors $\Psi_{r,s}^\pm(t)$ (for the relevant coefficients of the terms corresponding to the ordering kernel) based on explicit examples and also theoretical arguments. From these expressions one deduces that the parametrised vectors $\Psi_{r,s}^\pm(t)$ never vanish identically unlike the uncharged singular vectors of the Neveu-Schwarz $N = 2$ algebra at the degenerate points.

7 All twisted $N = 2$ singular vectors.

We have learned in section 2 that every positive parity singular vector $\Psi_{r,s}^+$, corresponding to an operator $(a, b)_l^+$, implies the existence of two singular vectors with negative parity at the same level l , corresponding to the operators $G(a, b)_l^-$ and $(a, b)_{G,l}^-$ (and the other way around for $\Psi_{r,s}^-$). These two negative (or positive) parity singular vectors may or may not be linearly independent. If they were always linearly independent, then whenever a singular vector exists we would find a two-dimensional singular space with the opposite parity. However, as was pointed out, the fact that $P(0) = 1$ in the determinant formula, given by Eq. (7), points towards the existence of, generically, one singular vector of each parity at the same level in the same Verma module $\mathcal{V}_{\Delta_{r,s}}$. Nevertheless, as we will see, even in the case where $G(a, b)_l^-$ and $(a, b)_{G,l}^-$ (or $G(a, b)_l^+$ and $(a, b)_{G,l}^+$) are proportional the maximal singular dimension may still be 2. In order to analyse this case in more detail we will use the expressions given by Eqs. (55)-(58) for $G(a, b)_l^-$, $(a, b)_{G,l}^-$, $G(a, b)_l^+$ and $(a, b)_{G,l}^+$.

Requiring that the two negative parity singular vectors are proportional, and using the explicit expressions for $\Delta_{r,s}(t)$, given by Eqs. (59), leads to two solutions: $\frac{b}{a} = 4\frac{rt}{rt+s}$ or $\frac{b}{a} = 4\frac{s}{rt+s}$. Note that under the proportionality assumption, the case $a = 0$ also implies $b = 0$ and thus $(a, b)_l^+ \equiv 0$, which we want to exclude. Therefore, if either of the two conditions on the ratio $\frac{b}{a}$ is satisfied, then the two negative parity singular vectors derived from a positive parity singular vector would be proportional and thus the corresponding singular dimension may well only be 1. Surprisingly enough, all examples found so far satisfy always the first of these two conditions. These examples include the singular vectors at levels $\frac{1}{2}$, 1 and $\frac{3}{2}$ given in section 6 as well as singular vectors at levels 2 and $\frac{5}{2}$. We believe that there must be a deep reason behind this result, i.e. underlying the fact that all the singular vectors we have computed satisfy the same condition (out of two possible ones) that prevents a positive parity singular vector to impose the existence of a two-dimensional singular space of negative parity. For this reason we conjecture that the positive parity singular vectors $\Psi_{r,s}^+ = (a, b)_{\frac{rs}{2}}^+ |\Delta_{r,s}\rangle$ follow the ratio $\frac{b}{a} = 4\frac{rt}{rt+s}$. From this we can easily derive the corresponding ratio $\frac{b'}{a'}$ for the negative parity singular vectors $\Psi_{r,s}^- = (a', b')_{\frac{rs}{2}}^- |\Delta_{r,s}\rangle$. These results allow us to conjecture the following expressions for the twisted $N = 2$ singular vectors.

Conjecture 7.A *The singular vectors $\Psi_{r,s}^\pm$ in the Verma modules $\mathcal{V}_{\Delta_{r,s}(t)}$ are given by:*

$$\Psi_{r,s}^+(t) = (rt + s, 4rt)_{\frac{rs}{2}}^+ |\Delta_{r,s}(t)\rangle, \quad (69)$$

$$\Psi_{r,s}^-(t) = \left(-\frac{2}{r}, rt + s\right)_{\frac{rs}{2}}^- |\Delta_{r,s}(t)\rangle. \quad (70)$$

Also for the negative parity singular vectors $\Psi_{r,s}^-$, the last two equations of Eqs. (55)-(58) both lead to $\Psi_{r,s}^+$ rather than to a pair of different singular vectors. The only exceptional cases occur when $\Psi_{r,s}^\pm$ or $\mathcal{V}_\Delta$ become G -closed and consequently some of these vectors vanish. This will be considered in the following sections.

Using the superconformal fusion rules it may be possible to derive explicit expressions for all the coefficients of the singular vectors Eqs. (69)-(70) in the spirit of Bauer, di Francesco, Itzykson, and Zuber^{4,3}, as it was possible for the Neveu-Schwarz $N = 2$ case¹⁰. This could also lead to a proof of conjecture 7.A. For this purpose it would be necessary to develop¹⁵ first a twisted superfield formalism following the approach of Ref. 13 for the Neveu-Schwarz algebras.

Based on conjecture 7.A it is easy to compute the Verma modules with degenerate singular vectors, i.e. with singular spaces of dimension 2. For a given level $l \in \mathbb{N}_0^{1/2}$ there are normally multiple solutions to the number theoretical problem $rs = 2l$, $r, s \in \mathbb{N}$, s odd. It happens the first time for level $l = \frac{3}{2}$ that we find multiple solutions $r = 1, s = 3$ and $r = 3, s = 1$. In the general case one factorises the integer $2l$ in its prime factors

$$2l = 2^n \prod_{i=1}^P p_i^{n_i}, \quad (71)$$

with $p_i > 2$ distinct primes and $n \in \mathbb{N}$, $n_i \in \mathbb{N}_0$. The solutions to $rs = 2l$, $r, s \in \mathbb{N}$, s odd are then given by

$$\begin{aligned} r_\pi &= 2^n \prod_{i=1}^P p_i^{k_i}, \\ s_\pi &= \prod_{i=1}^P p_i^{n_i - k_i}, \end{aligned} \quad (72)$$

for each P -tuple $\pi = (k_1, k_2, \dots, k_P) \in \mathbb{N}_0^P$, with $k_i \leq n_i, \forall i = 1, \dots, P$. For example, at level $l = 630$ we find the prime factorisation $2l = 2^2 \times 3^2 \times 5 \times 7$. Hence, we find 12 3-tuples $(k_1, k_2, k_3) \in \mathbb{N}_0^3$ with $k_1 \leq 2$ and $k_2, k_3 \leq 1$. We should note that for each level $l \in \mathbb{N}^{1/2}$ there exists at least one solution: $r = 2l, s = 1$. The number of solutions to $rs = 2l$, $r, s \in \mathbb{N}$, s odd is thus determined by the number of P -tuples $\pi = (k_1, \dots, k_P)$ with $\pi \in \mathbb{N}_0^P$ and $\pi \leq (n_1, \dots, n_P)$ for the prime factorisation of $2l$. Hence, there are^k $\Pi_l = \prod_{i=1}^P (n_i + 1)$ solutions to $rs = 2l$, $r, s \in \mathbb{N}$, s odd.

Due to our conjecture 7.A we have generically only one-dimensional singular spaces defined by the singular vectors $\Psi_{r,s}^+$ (or $\Psi_{r,s}^-$) at level $l = \frac{rs}{2}$ for all Π_l solutions of r and s . That is, these singular vectors with the same parity and level are located in different Verma modules. The most interesting Verma modules are those, where some of these Π_l solutions intersect and may hence lead to two-dimensional singular spaces provided the corresponding singular vectors are not proportional. Using conjecture 7.A we shall now investigate such cases of degeneration. Let us note again that the singular dimension can not be bigger than 2 even though a priori we would expect that even more than 2 solutions π_i could intersect.

^kWe define the empty product as $\prod_{i=1}^0 = 1$.

Let us analyse the intersections of the conformal weights $\Delta_{r,s}(t)$ of the Π_l solutions to $rs = 2l$, $r, s \in \mathbb{N}$, s odd. For fixed $l \in \mathbb{N}^{1/2}$ we find the prime factorisation as given above. Let us now take two P -tuples π^1 and π^2 ($\pi^1 \neq \pi^2$) and let us assume that the corresponding conformal weights intersect: $\Delta_{r_{\pi^1}, s_{\pi^1}} = \Delta_{r_{\pi^2}, s_{\pi^2}}$. Applying EQ. (59) one gets straightforwardly that this is the case if and only if

$$\left(2^n \prod_{i=1}^P p_i^{k_i^1} t + \prod_{i=1}^P p_i^{n_i - k_i^1} \right)^2 = \left(2^n \prod_{i=1}^P p_i^{k_i^2} t + \prod_{i=1}^P p_i^{n_i - k_i^2} \right)^2. \quad (73)$$

This easily rearranges to two sign-symmetric solutions for the parameter t of the central extension c :

$$t = \pm \frac{1}{2^n} \prod_{i=1}^P p_i^{n_i - k_i^1 - k_i^2}. \quad (74)$$

Again, let us consider the example of level $l = 630$ and consider the two 3-tuples $\pi^1 = (1, 1, 1)$ and $\pi^2 = (2, 0, 1)$ for the order $(3, 5, 7)$ of the relevant prime factors. EQ. (74) tells us that the conformal weights $\Delta_{r_{\pi^1}, s_{\pi^1}}$ and $\Delta_{r_{\pi^2}, s_{\pi^2}}$ have exactly 2 intersection points, namely $t = \pm \frac{1}{2^2 \times 3 \times 7} = \pm \frac{1}{84}$, where the two singular vectors $\Psi_{r_{\pi^1}, s_{\pi^1}}^+$ and $\Psi_{r_{\pi^2}, s_{\pi^2}}^+$ both exist. The key question will now be whether these singular vectors are actually different and hence define a two-dimensional singular space, or if they are proportional and thus lead to a one-dimensional singular space instead.

For this purpose, let us again take two different P -tuples π^1 and π^2 . The corresponding conformal weights intersect for $t = \pm \frac{1}{2^n} \prod_{i=1}^P p_i^{n_i - k_i^1 - k_i^2}$. Using conjecture 7.A we obtain the corresponding singular vectors $\Psi_{r_{\pi^j}, s_{\pi^j}}^+$, $j = 1, 2$ as:

$$\Psi_{r_{\pi^j}, s_{\pi^j}}^+ = (r_{\pi^j} t + s_{\pi^j}, 4r_{\pi^j} t)_l^+ \left| \Delta_{r_{\pi^j}, s_{\pi^j}} \right\rangle, j = 1, 2. \quad (75)$$

Inserting the expressions EQS. (72) for r_{π^j} and s_{π^j} and replacing t by EQ. (74) we hence obtain

$$\Psi_{r_{\pi^j}, s_{\pi^j}}^+ = \pm \left(\prod_{i=1}^P p_i^{n_i - k_i^{\bar{j}}} \pm \prod_{i=1}^P p_i^{n_i - k_i^j}, 4 \prod_{i=1}^P p_i^{n_i - k_i^{\bar{j}}} \right)_l^+ \left| \Delta_{r_{\pi^j}, s_{\pi^j}} \right\rangle, j = 1, 2, \quad (76)$$

with $\bar{1} = 2$ and $\bar{2} = 1$. The determinant of the coefficients shows that $\Psi_{r_{\pi^1}, s_{\pi^1}}^+$ and $\Psi_{r_{\pi^2}, s_{\pi^2}}^+$ are linearly dependent if and only if

$$\pm 4 \prod_{i=1}^P [p_i^{n_i - k_i^1}]^2 \mp 4 \prod_{i=1}^P [p_i^{n_i - k_i^2}]^2 = 0, \quad (77)$$

$$\Leftrightarrow \prod_{i=1}^P p_i^{n_i - k_i^1} = \prod_{i=1}^P p_i^{n_i - k_i^2}, \quad (78)$$

$$\Leftrightarrow \pi^1 = \pi^2, \quad (79)$$

in contradiction with our original assumption. Hence, the vectors $\Psi_{r_{\pi^1}, s_{\pi^1}}^+$ and $\Psi_{r_{\pi^2}, s_{\pi^2}}^+$ are linearly independent for $\pi^1 \neq \pi^2$. Similar arguments also show that $\Psi_{r_{\pi^1}, s_{\pi^1}}^-$ and $\Psi_{r_{\pi^2}, s_{\pi^2}}^-$ are linearly independent for $\pi^1 \neq \pi^2$ (at exactly the same values for t).

An important observation now is the following. On the one hand, we have proven that the maximal singular dimension is 2 and, on the other, we have shown that for different P -tuples π_1

and π_2 the corresponding singular vectors are always linearly independent at the intersection points (i.e. in the same Verma module). Therefore, since for high levels l there are many different P -tuples, the question arises whether or not it is possible that the conformal weights of more than two P -tuples can intersect at the same value t , obtaining a contradiction to either of our results. That this is not the case is shown in the following consideration. The conformal weights of π_1 and π_2 ($\pi_1 \neq \pi_2$) intersect at $t = \pm \frac{1}{2^n} \prod_{i=1}^P p_i^{n_i - k_i^1 - k_i^2}$. In the same way the conformal weights of π_2 and π_3 ($\pi_2 \neq \pi_3$) intersect at $\bar{t} = \pm \frac{1}{2^n} \prod_{i=1}^P p_i^{n_i - k_i^2 - k_i^3}$. Assuming $t = \bar{t}$ is equivalent to

$$\prod_{i=1}^P p_i^{n_i - k_i^1 - k_i^2} = \prod_{i=1}^P p_i^{n_i - k_i^2 - k_i^3}, \quad (80)$$

and thus $\prod_{i=1}^P p_i^{k_i^1} = \prod_{i=1}^P p_i^{k_i^3}$ results in $\pi_1 = \pi_3$ due to the fact that the numbers p_i are distinct primes. Therefore, there are no values of t for which more than two conformal weights Δ_{r_π, s_π} intersect for different P -tuples π . We should stress again, that these results are based on the conjecture 7.A which seems to be a very reliable conjecture. We summarise our results in the following theorem.

Theorem 7.B *Let us fix a level $l \in \mathbb{N}^{1/2}$. Based on the conjecture 7.A one finds that at the level $l = \frac{rs}{2}$ the singular vectors $\Psi_{r,s}^\pm$ define generically one-dimensional singular spaces. The only exceptions, where degenerate singular spaces of dimension 2 arise, occur necessarily for the cases when two conformal weights $\Delta_{r_1, s_1}(t)$ and $\Delta_{r_2, s_2}(t)$ intersect. For each two different pairs (r_1, s_1) and (r_2, s_2) with $l = \frac{r_1 s_1}{2} = \frac{r_2 s_2}{2}$, $r_1, r_2, s_1, s_2 \in \mathbb{N}$, s_1, s_2 odd, there are two values of t where such an intersection happens. These values are $t = \pm \frac{1}{2^n} \prod_{i=1}^P p_i^{n_i - k_i^1 - k_i^2}$ and they are real and rational in all cases. However, the conformal weights of three such pairs do not intersect for any common values of t .*

8 G -closed Verma modules.

We will now consider G -closed Verma modules $\mathcal{V}_{\frac{c}{24}}^G$ and discuss the relation between the singular vectors in these (incomplete) Verma modules and the singular vectors in complete Verma modules $\mathcal{V}_\Delta$ with $\Delta = \frac{c}{24}$. For this value the Verma module $\mathcal{V}_{\frac{c}{24}}$ contains a singular vector $\Psi_0^- = G_0 | \frac{c}{24} \rangle$ at level 0 with parity $-$. In order to obtain irreducible representations one considers the quotient space

$$\mathcal{V}_{\frac{c}{24}}^G = \frac{\mathcal{V}_{\frac{c}{24}}}{U(J_2)\Psi_0^-}, \quad (81)$$

which is precisely the G -closed Verma module $\mathcal{V}_{\frac{c}{24}}^G$. In fact, this quotient module is only one among the quotient modules. We could equally well consider quotients with respect to modules generated by singular vectors $\Psi_{r,s}^\pm$ at levels $\frac{rs}{2}$. The reason why we are especially interested in $\mathcal{V}_{\frac{c}{24}}^G$ is twofold. First, for physical applications it could be relevant if a highest weight vector (or a singular vector) is G -closed, which makes $\mathcal{V}_{\frac{c}{24}}^G$ special compared to other quotient spaces. Secondly, the Verma module $\mathcal{V}_{\frac{c}{24}}$ is easier to construct by explicit calculations than most Verma modules with singular vectors. Nevertheless, we aim also at obtaining information on the structure of quotient modules that is valid not only for $\mathcal{V}_{\frac{c}{24}}^G$ but also for quotient modules of similar type.

Via the canonical map defined for quotient spaces, singular vectors from $\mathcal{V}_{\frac{c}{24}}$ are either also singular in $\mathcal{V}_{\frac{c}{24}}^G$ or they are trivial (they ‘go away’). The latter happens if and only if the singular vector in question is a descendant vector of the singular vector Ψ_0^- in $\mathcal{V}_{\frac{c}{24}}$. Thus, singular vectors of $\mathcal{V}_{\frac{c}{24}}^G$ can either be inherited from $\mathcal{V}_{\frac{c}{24}}$ via the canonical quotient map or they appear for the first time in the quotient $\mathcal{V}_{\frac{c}{24}}^G$ as vectors that were not singular in $\mathcal{V}_{\frac{c}{24}}$. In the latter case these singular vectors of $\mathcal{V}_{\frac{c}{24}}^G$ are called subsingular vectors in $\mathcal{V}_{\frac{c}{24}}$. The Virasoro algebra does not allow any subsingular vectors which is a consequence of the proof of Feigin and Fuchs¹⁷, neither do the $N = 1$ superconformal algebras². Surprisingly enough, the $N = 2$ superconformal algebras contain subsingular vectors which have first been discovered in Refs. 20, 21 for the Neveu-Schwarz, the Ramond, and the topological $N = 2$ algebras. So far, nothing was known about the existence of subsingular vectors in the twisted $N = 2$ case. Due to the very different structure of this particular $N = 2$ algebra it was hard to say what one *a priori* should expect. However, we will show in this section that the twisted $N = 2$ superconformal algebra also contains subsingular vectors.

In section 5 we have found that the singular dimensions for G -closed Verma modules are just 1. Therefore, there are no degenerate singular vectors in G -closed Verma modules. A singular vector in $\mathcal{V}_{\frac{c}{24}}^G$ at level l can therefore be identified by the coefficient of only one particular term (being zero or non-zero). The ordering kernel in TAB. c tells us that this particular term is $T_{-\frac{1}{2}}^{2l}$ for the positive parity case and $T_{-\frac{1}{2}}^{2l-1}G_{-\frac{1}{2}}$ for the negative parity one. We can thus give the following theorem.

Theorem 8.A *Two singular vectors in the G -closed Verma module $\mathcal{V}_{\frac{c}{24}}^G$ at the same level and with the same parity are always proportional. If a vector satisfies the highest weight conditions at level l , with parity $+$ or parity $-$, and its coefficient for the term $T_{-\frac{1}{2}}^{2l}$ or $T_{-\frac{1}{2}}^{2l-1}G_{-\frac{1}{2}}$ vanishes, respectively, then this vector is trivial.*

Before looking at explicit examples of singular vectors in $\mathcal{V}_{\frac{c}{24}}^G$ we should first analyse whether or not the existing singular vectors $\Psi_{r,s}^\pm$ in $\mathcal{V}_{\frac{c}{24}}$ are descendants of Ψ_0^- . Requiring that $\Delta_{r,s}(t^G) = \frac{c}{24}$ in Eq. (59) one identifies easily the values t^G of t for which $\Psi_{r,s}^\pm$ exists in $\mathcal{V}_{\frac{c}{24}}$ at level $\frac{rs}{2}$:

$$t_{r,s}^G = -\frac{s}{r}, \quad (82)$$

for $r, s \in \mathbb{N}$, s odd. We observe that if $\Psi_{r,s}^\pm \in \mathcal{V}_{\frac{c}{24}}$, which happens only for the values $t_{r,s}^G$, then one finds also the singular vectors $\Psi_{nr,ns}^\pm \in \mathcal{V}_{\frac{c}{24}}$ at levels $\frac{n^2 rs}{2}$ for any positive integer $n \in \mathbb{N}$. Conjecture 7.A gives us the explicit form of the vectors $\Psi_{r,s}^\pm$. From $\Psi_{r,s}^+ = (rt + s, 4rt)_{\frac{rs}{2}}^+ | \frac{c}{24} \rangle$ one obtains for the values of $t_{r,s}^G$ that $\Psi_{r,s}^+(t_{r,s}^G) = (0, -4s)_{\frac{rs}{2}}^+ | \frac{r-s}{8r} \rangle$, and similarly $\Psi_{r,s}^- = (-\frac{2}{r}, rt + s)_{\frac{rs}{2}}^- | \frac{c}{24} \rangle$ leads to $\Psi_{r,s}^-(t_{r,s}^G) = (-\frac{2}{r}, 0)_{\frac{rs}{2}}^- | \frac{r-s}{8r} \rangle$. As the coefficient of $T_{-\frac{1}{2}}^{rs}$ for the positive parity case and the coefficient of $T_{-\frac{1}{2}}^{rs-1}G_{-\frac{1}{2}}$ in the negative parity case vanish, due to theorem 8.A we hence find that the singular vectors $\Psi_{r,s}^\pm(t_{r,s}^G)$ vanish in the quotient space $\mathcal{V}_{\frac{c}{24}}^G$. Therefore the singular vectors $\Psi_{r,s}^\pm(t_{r,s}^G)$ are descendant vectors of $\Psi_0^-(t_{r,s}^G)$. Consequently¹ all the singular vectors that can be found in $\mathcal{V}_{\frac{c}{24}}^G$ are subsingular in $\mathcal{V}_{\frac{c}{24}}$.

¹Strictly speaking, we have to assume that there are no isolated singular vectors for the twisted $N = 2$ superconformal algebra. We have not been able to compute any isolated singular vectors neither have they ever appeared for any other chiral algebra.

By explicit computation one easily finds that singular vectors $\Upsilon_{r,s}^\pm$ do appear in $\mathcal{V}_{\frac{r-s}{8r}}^G$ at exactly the levels $\frac{rs}{2}$ where the singular vectors $\Psi_{r,s}^\pm$ disappear in the quotient kernel. (The reasons for this will be shown in a future publication¹⁵). In addition these are the only singular vectors appearing in $\mathcal{V}_{\frac{c}{24}}^G$. The vectors $\Upsilon_{r,s}^\pm$ are hence subsingular vectors in the complete Verma modules $\mathcal{V}_{\frac{r-s}{8r}}$. This has been verified for levels $\frac{1}{2}$, 1, and $\frac{3}{2}$. The explicit results are given in appendix B.

We conclude this section with another interesting fact about the G -closed Verma modules. As we know, the singular dimension for G -closed Verma modules is only 1. One may wonder what happens in the cases where $\mathcal{V}_{\frac{c}{24}}$ has a two-dimensional singular space. First of all, we have just shown that if this happens then - due to the nature of the singular vectors $\Psi_{r,s}^\pm$ - the whole two-dimensional singular space would vanish in the quotient space $\mathcal{V}_{\frac{c}{24}}^G$ as it is descendant of Ψ_0^- . Nevertheless, one could argue that for these cases the corresponding subsingular vectors that appear in a one to one correspondence at the same levels than the singular vectors that disappear, may span a two-dimensional subsingular space. However, Verma modules $\mathcal{V}_\Delta$ that contain degenerate singular vectors never satisfy $\Delta = \frac{c}{24}$ and therefore this question is redundant. Namely, if we use in $t = -\frac{s}{r}$ the expressions of EQS. (72) for r and s and require that $t = \pm \frac{1}{2^n} \prod_{i=1}^P p_i^{n_i - k_i^1 - k_i^2}$ in the case of degeneracy for two P -tuples π^1 and π^2 , one easily finds

$$\pm \frac{1}{2^n} \prod_{i=1}^P p_i^{n_i - k_i^1 - k_i^2} = -\frac{1}{2^n} \prod_{i=1}^P p_i^{n_i - 2k_i^1} . \quad (83)$$

Taking into account that the p_i are distinct primes then the only solution to Eq. (83) is, however, $\pi^1 = \pi^2$, which shows that the modules $\mathcal{V}_\Delta$ with $\Delta = \frac{c}{24}$ never have two-dimensional singular spaces.

9 G -closed singular vectors.

In this last section we will analyse G -closed singular vectors in a very similar way as we have analysed G -closed Verma modules in the previous section. Due to the constraints on the conformal weights, there are no G -closed singular vectors in G -closed Verma modules, as mentioned already earlier. Therefore, we take $\Psi_{r,s}^\pm(t) \in \mathcal{V}_{\Delta_{r,s}(t)}$ and impose the conformal weight constraint $\Delta_{r,s} + \frac{rs}{2} = \frac{c}{24}$ which is satisfied by G -closed singular vectors. The condition $\Delta_{r,s} + \frac{rs}{2} = \frac{c}{24}$ has the unique solution

$${}^G t_{r,s} = \frac{s}{r} . \quad (84)$$

Let us note the similarity between ${}^G t_{r,s}$ and $t_{r,s}^G$ of the previous section.

According to the result that the singular dimension for G -closed singular vectors is 1, the ordering kernel being given in table TAB. b, a G -closed singular vector $\Psi_l^{\pm G}$ at level $l = \frac{c}{24} - \Delta$ can be identified by the coefficient of $T_{-\frac{1}{2}}^{2l}$ or $T_{-\frac{1}{2}}^{2l} G_0$ (being zero or non-zero), depending on the parity. We therefore give the following theorem.

Theorem 9.A *Two G -closed singular vectors in the same Verma module $\mathcal{V}_\Delta$, at the same level $l = \frac{c}{24} - \Delta$ and with the same parity are always proportional. If a G -closed vector satisfies the highest weight conditions at level $l = \frac{c}{24} - \Delta$ and parity +, or parity -, and its coefficient for the term $T_{-\frac{1}{2}}^{2l}$ or $T_{-\frac{1}{2}}^{2l} G_0$ vanishes, respectively, then this vector is trivial.*

Using the expressions for $G(a, b)_l^\mp$ given by Eqs. (55)-(58), we obtain for the vector $G_0\Psi_{r,s}^+ = G(rt + s, 4rt)_{\frac{rs}{2}}^- |\Delta_{r,s}\rangle$, in the case $t = {}^G t_{r,s}$, the result $G_0\Psi_{r,s}^+ = (0, 0)_{\frac{rs}{2}}^- |\Delta_{r,s}\rangle$. Since this vector satisfies the highest weight conditions but has vanishing coefficient for the term $T_{-\frac{1}{2}}^{rs} G_0$, we find that $G_0\Psi_{r,s}^+ \equiv 0$. Thus the singular vectors $\Psi_{r,s}^+$ are G -closed for $t = {}^G t_{r,s} = \frac{s}{r}$. Similarly, we obtain for $G_0\Psi_{r,s}^- = G(-\frac{2}{r}, rt + s)_{\frac{rs}{2}}^+ |\Delta_{r,s}\rangle$, in the case $t = {}^G t_{r,s}$, $G_0\Psi_{r,s}^- = (0, 0)_{\frac{rs}{2}}^+ |\Delta_{r,s}\rangle$ and hence the singular vectors $\Psi_{r,s}^-$ are also G -closed for $t = {}^G t_{r,s} = \frac{s}{r}$. It is surprising the fact that for $t = {}^G t_{r,s}$ all the singular vectors become G -closed (this is not required by first principles).

Theorem 9.B *The singular vectors $\Psi_{r,s}^\pm \in \mathcal{V}_{\Delta_{r,s}}$ become G -closed exactly for the values of the central parameter $t = {}^G t_{r,s} = \frac{s}{r}$, $r, s \in \mathbb{N}$, s odd.*

Like in the previous section, let us finally check that the cases where $\Psi_{r,s}^\pm \in \mathcal{V}_{\Delta_{r,s}}$ become G -closed cannot be degenerate cases, as suggested by the results of table TAB. b. We set $t = \frac{s}{r}$ and use again the expressions of Eqs. (72) for r and s . Requiring that $t = \pm \frac{1}{2^n} \prod_{i=1}^P p_i^{n_i - k_i^1 - k_i^2}$ in the case of degeneracy for two different P -tuples π^1 and π^2 we obtain

$$\pm \frac{1}{2^n} \prod_{i=1}^P p_i^{n_i - k_i^1 - k_i^2} = + \frac{1}{2^n} \prod_{i=1}^P p_i^{n_i - 2k_i^1}. \quad (85)$$

Again, the only solution to Eq. (85) is $\pi^1 = \pi^2$, in contradiction with the assumption that π^1 and π^2 are distinct. This shows that the degenerate cases do not correspond to G -closed singular vectors, i.e. two-dimensional singular spaces are never G -closed.

10 Conclusions and prospects

In this paper we have applied the method of adapted orderings to the twisted $N = 2$ superconformal algebra. This shows, again, that this method is a powerful tool that can easily be applied to a large classes of algebras. The way it works is rather simple and it is very flexible in its use. Whereas the application of this method to the topological, Neveu-Schwarz and Ramond $N = 2$ algebras in Ref. 16 assigned a special rôle to the Virasoro generator L_{-1} , for the twisted $N = 2$ algebra this special rôle has been taken over by the $U(1)$ current generator $T_{-\frac{1}{2}}$. Another remarkable difference compared to our earlier applications of this method is that in the twisted $N = 2$ case the fermionic field modes need to be mixed in the adapted ordering rather than being kept separate.

As a result we have derived the singular dimensions and the ordering kernels for the twisted $N = 2$ algebra. This reveals that also the twisted $N = 2$ algebra contains degenerate singular spaces with dimension 2 in complete Verma modules, as it is the case for the three isomorphic $N = 2$ algebras. On the other hand, G -closed Verma modules have singular dimension 1 and therefore do not allow any degenerate singular spaces.

Based on the ordering kernel coefficients we have conjectured general expressions for the relevant terms of all (primitive) singular vectors. These expressions lead to the identification of all degenerate cases with singular dimension 2 and also to the identification of all G -closed singular vectors. These expressions also lead to the discovery of subsingular vectors for the twisted $N = 2$ superconformal algebra. As in many examples for the other $N = 2$ superconformal algebras, these subsingular vectors appear in relation to additional vanishing conditions on a singular vector. This relation can be generally proven and will be given in a forthcoming paper¹⁵. We have seen that ordering kernel coefficients play a crucial rôle in this matter.

We have also computed multiplication rules for singular vector operators, which sets the foundation for the study of the twisted $N = 2$ embedding diagrams which will be investigated in a future publication.

The twisted $N = 2$ superconformal algebra has hence proven to have a rather rich structure that contains many features that were also discovered for the three isomorphic $N = 2$ superconformal algebras. Nevertheless, the way these features appear in the case of the twisted $N = 2$ algebra is quite different from the way they appear for the other $N = 2$ algebras. For example, the degenerate singular spaces are produced simply by the intersection of two primitive singular vectors. For no other $N = 2$ algebra primitive singular vectors at the same level and with the same charge are linearly independent. Furthermore, the discovery of degenerate singular vectors in the twisted $N = 2$ case is rather surprising as the underlying parameter space is just one-dimensional. The twisted $N = 2$ superconformal algebra hence turns out to be quite distinct from all other superconformal algebras analysed so far in the literature. This raises the need for the analysis of twisted superconformal algebras for higher N - a very interesting question already because the periodicity conditions of more than 2 fermionic fields can be mixed in different ways. The concept of adapted orderings should apply also in these cases.

A Examples of negative parity singular vectors

For completeness we shall also give all singular vectors with negative parity at level $\frac{1}{2}$, 1, and $\frac{3}{2}$. They were found by explicit computer calculations²⁹.

Level $\frac{1}{2}$:

$$\Psi_{1,1}^-(t) = \left\{ -2T_{-\frac{1}{2}}G_0 + (t+1)G_{-\frac{1}{2}} \right\} \left| -\frac{t+1}{8t} \right\rangle. \quad (86)$$

Level 1:

$$\begin{aligned} \Psi_{2,1}^-(t) = & \left\{ -T_{-\frac{1}{2}}^2G_0 + (2t+1)T_{-\frac{1}{2}}G_{-\frac{1}{2}} - 2tL_{-1}G_0 \right. \\ & \left. - \frac{t}{2}(2t+3)G_{-1} \right\} \left| -\frac{3t^2+3t+1}{8t} \right\rangle. \end{aligned} \quad (87)$$

Level $\frac{3}{2}$ has two different singular vectors:

$$\begin{aligned} \Psi_{3,1}^-(t) = & \left\{ -\frac{2}{3}T_{-\frac{1}{2}}^3G_0 + (3t+1)T_{-\frac{1}{2}}^2G_{-\frac{1}{2}} - 4tL_{-1}T_{-\frac{1}{2}}G_0 \right. \\ & - \frac{1}{3}(12t^2+13t+3)G_{-1}T_{-\frac{1}{2}} + \frac{2t}{3}(3t+1)L_{-1}G_{-\frac{1}{2}} - \frac{4t}{3}G_{-1}G_{-\frac{1}{2}}G_0 \\ & \left. - \frac{4t}{3}(t-1)T_{-\frac{3}{2}}G_0 + (2t^3 - \frac{4}{3}t^2 - \frac{8}{3}t - \frac{2}{3})G_{-\frac{3}{2}} \right\} \left| -\frac{8t^2+5t+1}{8t} \right\rangle, \end{aligned} \quad (88)$$

$$\begin{aligned} \Psi_{1,3}^-(t) = & \left\{ -2T_{-\frac{1}{2}}^3G_0 + (t+3)T_{-\frac{1}{2}}^2G_{-\frac{1}{2}} + 4tL_{-1}T_{-\frac{1}{2}}G_0 \right. \\ & - (t+3)G_{-1}T_{-\frac{1}{2}} - 2t(t+3)L_{-1}G_{-\frac{1}{2}} - 4tG_{-1}G_{-\frac{1}{2}}G_0 \\ & \left. + 8T_{-\frac{3}{2}}G_0 - 2(t+3)G_{-\frac{3}{2}} \right\} \left| -\frac{5t+9}{8t} \right\rangle. \end{aligned} \quad (89)$$

In the notation of definition 5.B we obtain:

$$\Psi_{1,1}^-(t) = (-2, t+1)_{\frac{1}{2}}^- \left| -\frac{t+1}{8t} \right\rangle, \quad (90)$$

$$\Psi_{2,1}^-(t) = (-1, 2t+1)_1^- \left| -\frac{3t^2+3t+1}{8t} \right\rangle, \quad (91)$$

$$\Psi_{3,1}^-(t) = \left(-\frac{2}{3}, 3t+1\right)_{\frac{3}{2}}^- \left| -\frac{8t^2+5t+1}{8t} \right\rangle, \quad (92)$$

$$\Psi_{1,3}^-(t) = (-2, t+3)_{\frac{3}{2}}^- \left| -\frac{5t+9}{8t} \right\rangle. \quad (93)$$

For $t = \pm 1$ we observe that $\Psi_{3,1}^-$ and $\Psi_{1,3}^-$ span a two-dimensional singular vector space, as it was the case for their positive parity partners.

$$\Psi_{3,1}^-(1) = \left(-\frac{2}{3}, 4\right)_{\frac{3}{2}}^- \left| -\frac{7}{4} \right\rangle, \quad (94)$$

$$\Psi_{1,3}^-(1) = (-2, 4)_{\frac{3}{2}}^- \left| -\frac{7}{4} \right\rangle, \quad (95)$$

$$\Psi_{3,1}^-(-1) = \left(-\frac{2}{3}, -2\right)_{\frac{3}{2}}^- \left| \frac{1}{2} \right\rangle, \quad (96)$$

$$\Psi_{1,3}^-(-1) = (-2, 2)_{\frac{3}{2}}^- \left| -\frac{1}{2} \right\rangle. \quad (97)$$

B Examples of singular vectors in G -closed Verma modules

As discussed in section 8, the singular vectors $\Psi_{r,s}^\pm \in \mathcal{V}_{\Delta_{r,s}}$, at levels $\frac{rs}{2}$, are always descendants of the level zero singular vector $\Psi_0^- = G_0 \left| \frac{c}{24} \right\rangle$ in the cases where $\Delta_{r,s}(t) = \frac{c}{24}$. This happens for the values $t = t_{r,s}^G = -\frac{s}{r}$, so that $\frac{c}{24} = \frac{r-s}{8r}$. Therefore, the singular vectors $\Psi_{r,s}^\pm$ are absent in the G -closed Verma modules $\mathcal{V}_{\frac{c}{24}}^G = \frac{\mathcal{V}_{\frac{c}{24}}}{G_0 \left| \frac{c}{24} \right\rangle}$. All singular vectors which we find in $\mathcal{V}_{\frac{c}{24}}^G$ are therefore subsingular vectors in $\mathcal{V}_{\frac{c}{24}}$. Surprisingly enough, the only singular vectors we have found in $\mathcal{V}_{\frac{c}{24}}^G$ are exactly at the levels and for the modules where we would have expected $\Psi_{r,s}^\pm$ to appear, if they were not trivial in the quotient module. That is, they are located at levels $\frac{rs}{2}$ in the Verma modules with $t = -\frac{s}{r}$. In the following we will give all the singular vectors in $\mathcal{V}_{\frac{c}{24}}^G$ for levels $\frac{1}{2}$, 1, and $\frac{3}{2}$. These vectors are therefore subsingular vectors in $\mathcal{V}_{\frac{c}{24}}$.

Level $\frac{1}{2}$ has singular vectors only for $t = -1$:

$$\Upsilon_{1,1}^+(-1) = T_{-\frac{1}{2}} |0\rangle^G, \quad (98)$$

$$\Upsilon_{1,1}^-(-1) = G_{-\frac{1}{2}} |0\rangle^G. \quad (99)$$

Level 1 has singular vectors only for $t = -\frac{1}{2}$:

$$\Upsilon_{2,1}^+(-\frac{1}{2}) = \left\{ T_{-\frac{1}{2}}^2 - L_{-1} \right\} \left| \frac{1}{16} \right\rangle^G, \quad (100)$$

$$\Upsilon_{2,1}^-(-\frac{1}{2}) = \left\{ 4T_{-\frac{1}{2}} G_{-\frac{1}{2}} - G_{-1} \right\} \left| \frac{1}{16} \right\rangle^G. \quad (101)$$

Level $\frac{3}{2}$ has singular vectors only for $t = -\frac{1}{3}$ and for $t = -3$:

$$\Upsilon_{3,1}^+(-\frac{1}{3}) = \left\{ 9T_{-\frac{1}{2}}^3 - 18L_{-1}T_{-\frac{1}{2}} - 6G_{-1}G_{-\frac{1}{2}} + 8T_{-\frac{3}{2}} \right\} \left| \frac{1}{12} \right\rangle^G, \quad (102)$$

$$\Upsilon_{3,1}^-(-\frac{1}{3}) = \left\{ 27T_{-\frac{1}{2}}^2G_{-\frac{1}{2}} - 15G_{-1}T_{-\frac{1}{2}} - 6L_{-1}G_{-\frac{1}{2}} - 10G_{-\frac{3}{2}} \right\} \left| \frac{1}{12} \right\rangle^G. \quad (103)$$

$$\Upsilon_{1,3}^+(-3) = \left\{ T_{-\frac{1}{2}}^3 + 6L_{-1}T_{-\frac{1}{2}} - 6G_{-1}G_{-\frac{1}{2}} - 4T_{-\frac{3}{2}} \right\} \left| -\frac{1}{4} \right\rangle^G, \quad (104)$$

$$\Upsilon_{1,3}^-(-3) = \left\{ T_{-\frac{1}{2}}^2G_{-\frac{1}{2}} - G_{-1}T_{-\frac{1}{2}} + 6L_{-1}G_{-\frac{1}{2}} - 2G_{-\frac{3}{2}} \right\} \left| -\frac{1}{4} \right\rangle^G. \quad (105)$$

If we replace $\left| \frac{r-s}{8r} \right\rangle^G$ by $\left| \frac{r-s}{8r} \right\rangle$ the resulting singular vectors are thus subsingular vectors in the Verma modules $\mathcal{V}_{\frac{r-s}{8r}}$.

The appearance of these subsingular vectors is very closely related with an additional vanishing condition for the singular vector $G_0 \left| \frac{c}{24} \right\rangle$. That such an additional vanishing condition results in an additional null vector which is most likely subsingular will be proven in a forthcoming publication¹⁵.

Acknowledgements

We are grateful to Adrian Kent for many discussions on the ordering method and to Victor Kac for his explanations on superconformal algebras. M.D. is indebted to the Deutsche Forschungsgemeinschaft (DFG) for financial support and to DAMTP for the hospitality.

References

- [1] M. Ademollo, L. Brink, A. d'Adda, R. d'Auria, E. Napolitano, S. Sciuto, E. del Giudice, P. di Vecchia, S. Ferrara, F. Gliozzi, R. Musto, and R. Pettorino. Supersymmetric strings and colour confinement. *Phys. Lett.*, B62:105, 1976.
- [2] A.B. Astashkevich. On the structure of Verma modules over Virasoro and Neveu-Schwarz algebras. *Commun. Math. Phys.*, 186:531, 1997.
- [3] M. Bauer, P. di Francesco, C. Itzykson, and J.-B. Zuber. Covariant differential equations and singular vectors in Virasoro representations. *Nucl. Phys.*, B362:515, 1991.
- [4] M. Bauer, P. di Francesco, C. Itzykson, and J.-B. Zuber. Singular vectors of the Virasoro algebra. *Phys. Lett.*, B260:323, 1991.
- [5] L. Benoit and Y. Saint-Aubin. Fusion and the Neveu-Schwarz singular vectors. *Lett. Math. Phys.*, 23:117, 1991.
- [6] L. Benoit and Y. Saint-Aubin. An explicit formula for some singular vectors of the Neveu-Schwarz algebra. *Int. J. Mod. Phys.*, A7:3023, 1992.
- [7] M. Bershadsky, W. Lerche, D. Nemeschansky and N.P. Warner. Extended $N = 2$ superconformal structure of gravity and W-gravity coupled to matter. *Nucl. Phys.*, B401:304, 1993.

- [8] W. Boucher, D. Friedan, and A. Kent. Determinant formulae and unitarity for the $N = 2$ superconformal algebras in two dimensions or exact results on string compactification. *Phys. Lett.*, B172:316, 1986.
- [9] R. Dijkgraaf, E. Verlinde, and H. Verlinde. Topological strings in $d < 1$. *Nucl. Phys.*, B352:59, 1991.
- [10] M. Dörrzapf. Singular vectors of the $N = 2$ superconformal algebra. *Int. J. Mod. Phys.*, A10:2143, 1995.
- [11] M. Dörrzapf. *Superconformal field theories and their representations*. PhD thesis, <http://www.damtp.cam.ac.uk/user/md131/research/thesis.html>, University of Cambridge, September 1995.
- [12] M. Dörrzapf. Analytic expressions for singular vectors of the $N = 2$ superconformal algebra. *Commun. Math. Phys.*, 180:195, 1996.
- [13] M. Dörrzapf. The definition of Neveu-Schwarz superconformal fields and uncharged superconformal transformations. *HUTP-97/A054 preprint, hep-th/9712197, to be published in Rev.Math. Phys.*, 1999.
- [14] M. Dörrzapf. The embedding structure of unitary $N = 2$ minimal models. *Nucl. Phys.*, B529:639, 1998.
- [15] M. Dörrzapf and B. Gato-Rivera. Work in progress.
- [16] M. Dörrzapf and B. Gato-Rivera. Singular dimensions of the $N = 2$ superconformal algebras. I. *IMAFF-FM-98/06, NIKHEF-98-022, HUTP-98/A021 preprint, hep-th/9807234*, 1998.
- [17] B.L. Feigin and D.B. Fuchs. *Representations of Lie groups and related topics*. A.M. Vershik and A.D. Zhelobenko eds. , Gordon & Breach, 1990.
- [18] D. Friedan, Z. Qiu, and S. Shenker. Superconformal invariance in two dimensions and the tricritical Ising model. *Phys. Lett.*, B151:37, 1985.
- [19] D. Fuchs. Singular vectors over the Virasoro algebra and extended Verma modules. *Adv. Sov. Math.*, 17:65, 1993.
- [20] B. Gato-Rivera and J.I. Rosado. Chiral determinant formula and subsingular vectors for the $N = 2$ superconformal algebras. *Nucl. Phys.*, B503:447, 1997.
- [21] B. Gato-Rivera and J.I. Rosado. Families of singular and subsingular vectors of the topological $N = 2$ superconformal algebra. *Nucl. Phys.*, B514:477, 1998.
- [22] B. Gato-Rivera and A.M. Semikhatov. $d \leq 1$ U $d \geq 25$ and W constraints from BRST-invariance in the $c \neq 3$ topological algebra. *Phys. Lett.*, B293:72, 1992.
- [23] P. Goddard, A. Kent, and D. Olive. Unitary representations of the Virasoro and super-Virasoro algebras. *Commun. Math. Phys.*, 103:105, 1986.
- [24] V.G. Kac. Lie superalgebras. *Adv. Math.*, 26:8, 1977.
- [25] V.G. Kac. Contravariant form for infinite-dimensional (Lie) algebras and superalgebras. *Lec. Notes in Phys.*, 94:441, 1979.
- [26] A. Kent. Singular vectors of the Virasoro algebra. *Phys. Lett.* B273:56, 1991.

- [27] E.B. Kiritsis. Character formulae and the structure of the representations of the $N = 1$, $N = 2$ superconformal algebras. *Int. J. Mod. Phys.*, A3:1871, 1988.
- [28] A. Meurman and A. Rocha-Caridi. Highest weight representations of the Neveu-Schwarz and Ramond algebras. *Commun. Math. Phys.*, 107:263, 1986.
- [29] Waterloo Maple Software. Programmed with Maple V, release 3, 1998.
- [30] G.M.T. Watts. Null vectors of the superconformal algebra: The Ramond sector. *Nucl. Phys.*, B407:213, 1993.